
### **Regional Seas and GEF-LME Linkages**

- The Global Environment Facility (GEF) is a funding agency assisting developing coastal countries to meet ecosystem-related targets. The GEF recommends the use of LMEs as the geographic focus for ecosystem-based assessments and management strategies.
- In 18 GEF-LME projects currently underway, project goals include the joint preparation of a Transboundary Diagnostic Analysis (TDA) and a Strategic Action Plan (SAP), used to prioritize Project actions.
- In a GEF-LME Project TDA, the countries bordering the LME prepare a document based on consensus that ranks coastal resource issues, identifies and prioritizes transboundary problems, analyzes socioeconomic impacts, outlines root causes and advances possible remedies
- On the basis of the TDA, the countries prepare a Strategic Action Plan (SAP). In the SAP, the countries propose to remedy the transboundary issues identified in the TDA and outline national and regional commitments to policy, legal and institutional reform.
- Countries follow Project goals and milestones leading towards an adaptive, ultimately self-financing, management regime for LMEs located within Regional Seas areas around the globe.
- The TDA and SAP processes for LMEs are consistent with the 2002 Johannesburg targets and Plan of Implementation (POI) of the World Summit on Sustainable Development (WSSD)

## The WSSD targets are useful criteria by which to gauge the progress of GEF-LME Projects

- Achievement of substantial reductions in land-based sources of pollution by 2006
- Introduction of an ecosystems approach to marine resource assessment and management by 2010
- Designation of a network of marine protected areas by 2012
- Maintenance and restoration of fish stocks to maximum sustainable yield levels by 2015



18 GEF-LME Projects in Regional Seas. GEF-LME Projects in Regional Seas already approved (red dots) or in the preparation stage (yellow dots), involve 121 countries of Africa, Asia, the Pacific, Latin America & the Caribbean, and Eastern Europe.

### For further information contact:



## NOAA — Large Marine Ecosystem Program Narragansett Laboratory, NOAA-NMFS

28 Tarzwell Drive, Narragansett, RI 02882, USA

Phone: +1 (401) 782-3211 FAX: +1 (401) 782-3201

Email: Kenneth.Sherman@NOAA.gov Website: www.edc.uri.edu/lme



### INEP — United Nations Environment Programme

Regional Seas Programme
PO Box 30552, Nairobi, Kenya
Regional Seas Website: www.unep.ch/regionalseas

GPA — Global Programme of Action for the Protection of the Marine

Environment from Land-Based Activities

PO Box 16227, 2500 BE, The Hague, Netherlands

Phone: +31 70 311 4460 FAX: +31 70 345 6648

Email: veerle.rs@unep.nl



### GEF — The Global Environment Facility

Website: www.gpa.unep.org

GEF Secretariat 1818 H Street, NW Washington, DC, USA

Phone: +1 (202) 473-0508 FAX: +1 (202) 522-3240/3245

Email: aduda@thegef.org Website: www.gefweb.org

### **Network of United Nations Organizations and NGOs participating in GEF-LME Projects**

### INTERNATIONAL PARTNERS

GEF — The Global Environment Facility

The World Bank

IOC — Intergovernmental Oceanographic Commission

UNEP — United Nations Environment Programme

Regional Seas Programme

Global International Waters Assessment (GIWA)

Global Programme of Action for the Protection of the Marine Environment from Land-Based Activities (GPA)

UNDP (United Nations Development Programme)

UNIDO (United Nations Industrial Development Organization)

FAO (Food and Agriculture Organization, Fisheries Division)

IUCN (International Union for the Conservation of Nature—World Conservation Union)

WWF (World Wildlife Fund)

### **Bibliographical references**

- Duda, A. M. and K. Sherman, 2002. A new imperative for improving management of large marine ecosystems. Ocean and Coastal Management 45:797-833.
- Koranteng, K. A., 2002. In J. McGlade, P. Cury, K. Koranteng, N. J. Hardman-Mountford, "The Gulf of Guinea Large Marine Ecosystem: Environmental Forcing and Sustainable Development of Marine Resources." Elsevier Science. 392p.
- Kroeze, C. and S. Seitzinger, 1998. Nitrogen inputs to rivers, estuaries and continental shelves and related nitrous oxide emissions in 1990 and 2050. Nutrient Cycling in Agroecosystems 52:195-212.
- Kumpf, H., K. Steidinger, and K. Sherman, 1999. "The Gulf of Mexico Large Marine Ecosystem: Assessment, Sustainability, and Management." Blackwell Science. 736p.
- Sherman, K. and A.M. Duda, 1999. An ecosystem approach to global assessment and management of coastal waters. Marine Ecology Progress Series. 190:271-287.








## **UNEP Regional Seas Programme**

Linked with

# Large Marine Ecosystems

**Assessment and Management** 



Regional Seas, West to East: North-East Pacific • South-East Pacific • Wider Caribbean • South-West Atlantic • West & Central Africa Mediterranean • Black Sea • Eastern Africa • Red Sea & Gulf of Aden • ROPME Sea Area • South Asian Seas • East Asian Seas North-West Pacific • South Pacific Independent Partners: Arctic • North-East Atlantic • Baltic Sea • Caspian Sea • Antarctic

A new partnership has been developed that links the coast and oceans activities of the United Nations Environment Programme (UNEP) and the US National Oceanographic and Atmospheric Administration (NOAA). The joint initiative incorporates NOAA's Large Marine Ecosystem (LME) assessment and management approach, using LMEs as operational/management units for translating the Regional Seas Programme into concrete actions. This will assist countries in Africa, Asia, Latin America & the Caribbean, and Eastern Europe to restore and sustain resources, coastal environments and linked watersheds.

- The UNEP Regional Seas Programme is focused on assisting countries in protecting the coastal and marine environment. Regional Seas promotes and advances coastal programmes and legal agreements that help countries move forward to control coastal degradation.
- ❖ In 2004, a new strategic initiative was adopted by the Regional Seas Programme. Its aim amongst others is to advance "the development of a common vision and integrated management, based on the ecosystem approach, of priorities and concerns related to the coastal and marine environment and its resources in Regional Seas Conventions and Action Plans."
- Based on this priority a new partnership has been developed that links the Regional Seas Programme and Large Marine Ecosystem (LME) Projects. It focuses on the assessment and management of LMEs located in Regional Seas areas.

## **Regional Seas Partners with GEF-LME Projects**

### UNEP

Since 1974, the United Nations Environment Programme (UNEP), through its Regional Seas Programme, has been engaged in assisting countries in protecting their marine environment. The UNEP-Regional Seas Programme has developed regional action-oriented programmes and legally-binding regional conventions for countries bordering the 18 Regional Seas. In 1982, UNEP began to address issues related to impacts on the marine environment from land-based activities. The response to intense pressures put on coastal systems was the 1995 Washington Declaration, by 108 governments and the European Commission, to adopt a Global Programme of Action (GPA) for the Protection of the Marine Environment from Land-based Activities. UNEP was tasked to lead the coordination effort and to establish a GPA Coordination Office.

## The GPA and the control of land-based sources of pollution

Some 80% of the pollution load in the oceans originates from land-based activities (municipal, industrial and agricultural wastes, run-off, and atmospheric deposition). These contaminants affect the most productive areas of the marine environment, including estuaries and near-shore coastal waters. The health, and in some cases the very survival, of coastal populations depend upon the health and well being of coastal systems such as estuaries and wetlands. To support the GPA activity, a UNEP/GPA office was established in The Hague, Netherlands. One strategic direction for GPA/Regional Seas in the coming decade is to promote ecosystem-based management.

### **UNEP's Regional Seas and LMEs**

Large Marine Ecosystems (LMEs) are regions of ocean space of about 200,000 km2 or greater. They encompass coastal areas from river basins and estuaries out seaward to the break or slope of the continental shelf (e.g. Yellow Sea LME), or out to the seaward extent of a

well-defined current system (e.g. Guinea Current LME). Some, like the Black Sea LME, are semi-enclosed geographical areas. LMEs are defined by ecological criteria including (1) bottom depth contours, (2) currents and water mass structure, (3) marine productivity, and (4) food webs. LMEs are located within the Regional Seas areas.

- LMEs annually produce 95% of the world's marine fish catch.
- SeaWiFS satellite data provide estimates on LME primary productivity.
- Assessments of the changing conditions in LMEs include fish and fisheries, and the influence of land-based sources of pollutants and disruption of the nitrogen cycle on the integrity and health of LMEs.

LMEs provide a flexible approach to ecosystem-based management by identifying driving forces of ecosystem change. Since 1984, NOAA has been supporting the development of LME assessment and management strategies, in partnership with the IOC, the IUCN and United Nations agencies. Twelve peer-reviewed volumes and case studies of LME management strategies have been published since 1986 (Duda and Sherman, 2002).

The Global Environment Facility (GEF) is a funding

### **GEF-LME Projects in Regional Seas areas**

agency that partners with national and international agencies to assist developing coastal countries in meeting the four ecosystem-related targets agreed upon at the 2002 World Summit on Sustainable Development (WSSD) held in Johannesburg. Since the early 1990s, developing countries have approached the GEF, and U.N. implementing agencies, for technical and scientific assistance in restoring and protecting their coastal and marine ecosystems. The GEF recommends the use of LMEs as the geographic focus for ecosystem-based strategies to reduce coastal pollution, restore damaged habitats, and recover depleted fisheries.

## **UNEP Regional Seas Programme Linked with the LME Approach**

The LME approach is a way forward for promoting ecosystem-based management of coastal and marine resources within a framework of sustainable development. Country-driven GEF-LME assessment and management projects are linked to the WSSD Plan of Implementation and to the UNEP Regional Seas Programme

- In 18 GEF-LME Projects (either approved or in the preparation stage), 121 countries are moving to meet ecosystem-related targets and to address overfishing, fishing down food webs, destruction of habitat and accelerated nitrogen export.
- In 14 Projects, 111 countries engaged in the Transboundary Diagnostic Analysis (TDA) process have already begun to scientifically characterize the LME, to identify the root causes of trends in LME biomass yields and the most pressing transboundary characteristics of coastal pollution, damaged habitats and depleted fish stocks, in order to prioritize these issues.
- 7 country-driven GEF-LME Projects are advancing to the drafting of the Strategic Action Plan (SAP), in which the countries commit to making institutional arrangements and taking policy actions, based on sound science, to address the issues identified in the TDA. The SAP addresses to correct institutional fragmentation, ecosystem assesment gaps, lack of cooperation and weak coastal policies and is signed by high-level government authorities of each participating country.
- In 13 LMEs out of 29 LME case studies, climate forcing was discovered to be the principal driver of change in biomass yield; in 14 LMEs, it was overfishing; and in one LME, eutrophication. For one LME, the results were inconclusive.
- The strategic framework for developing TDAs and SAPs is guided by the geographic area of LMEs and the application of the 5-module approach to LME assessment and management.

Table of country-driven GEF-LME Projects with Regional Seas that have established priorities through a Transboundary Diagnostic Analysis and a Strategic Action Programme planning process:

### **GEF-LME Projects Regional Seas Benguela Current (3 countries)** West and Central Africa **TDA and SAP** Red Sea (7 countries) **TDA and SAP** Red Sea and Gulf of Ader Mediterranean Sea (19 countries) **TDA and SAP** Mediterranean Sea Gulf of Guinea (6 countries) **TDA and SAP West and Central Africa TDA and SAP** Black Sea (6 countries) Black Sea **South China Sea (7 countries) TDA and SAP East Asian Seas TDA and SAP North-West Pacific** Yellow Sea (2 countries) **TDA and SAP Baltic Sea (9 countries) Baltic Sea** Bay of Bengal (8 countries) **TDA** and preliminary **SAP South Asian Seas Canary Current (7 countries) TDA** in preparation **West and Central Africa Humboldt Current (2 countries) TDA** in preparation **South-East Pacific** Guinea Current (16 countries) Expanding the SAP and TDA West and Central Africa **Gulf of Mexico (3 countries) TDA** in preparation Wider Caribbean Caribbean Sea (23 countries) **TDA** in preparation Wider Caribbean

LME 13.

LME 23.

LME 26...

LME 32...

LME 34.


LME 61.

LME 62 ..

LMEs 14,15,16,17..

## Regional Seas & Large Marine Ecosystems







Regional Seas, West to East: North-East Pacific • South-East Pacific • Wider Caribbean • South-West Atlantic • West & Central anean • Black Sea • Eastern Africa • Red Sea & Gulf of Aden • ROPME Sea Area • South Asian Seas • East Asian Seas North-West Pacific • South Pacific Independent Partners: Arctic • North-East Atlantic • Baltic Sea • Caspian Sea • Antarctic

### **64 LMEs of the World**

3 California Current 16 East Brazil Shelf 29 Benguela Current 41 East-Central Australian Shelf 54 Chukchi 4 Gulf of California 17 North Brazil Shelf 30 Agulhas Current 42 Southeast Australian Shelf 55 Beauford 55 Gulf of Mexico 18 West Greenland Shelf 31 Somali Coastal Current 43 Southwest Australian Shelf 56 East Sib Southeast U.S. Continental Shelf 19 East Greenland Shelf 32 Arabian Sea 44 West-Central Australian Shelf 57 Laptev Southast U.S. Continental Shelf 20 Barents Sea 33 Red Sea 45 Northwest Australian Shelf 58 Kara Sea Sotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland Southast Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic 11 Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Se	I East Bering Sea	14 Patagonian Shelf	27 Canary Current	40 Northeast Australian Shelf-	52 Okhotsk Sea
4 Gulf of California 17 North Brazil Shelf 30 Agulhas Current 42 Southeast Australian Shelf 55 Beauford 56 Gulf of Mexico 18 West Greenland Shelf 31 Somali Coastal Current 43 Southwest Australian Shelf 56 East Sib Southeast U.S. Continental Shelf 19 East Greenland Shelf 32 Arabian Sea 44 West-Central Australian Shelf 57 Laptev S Northeast U.S. Continental Shelf 20 Barents Sea 33 Red Sea 45 Northwest Australian Shelf 58 Kara Sea 8 Scotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland S Gulf of Thailand 47 East China Sea 60 Faroe Pl In Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic II Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Se	2 Gulf of Alaska	15 South Brazil Shelf	28 Guinea Current	Great Barrier Reef	53 West Bering Sea
5 Gulf of Mexico 18 West Greenland Shelf 31 Somali Coastal Current 43 Southwest Australian Shelf 56 East Sib Southeast U.S. Continental Shelf 19 East Greenland Shelf 32 Arabian Sea 44 West-Central Australian Shelf 57 Lapter S 7 Northeast U.S. Continental Shelf 20 Barents Sea 33 Red Sea 45 Northwest Australian Shelf 58 Kara Sea 8 Scotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland S 9 Newfoundland-Labrador Shelf 22 North Sea 35 Gulf of Thailand 47 East China Sea 60 Faroe Pl 10 Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic 11 Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Sea	3 California Current	16 East Brazil Shelf	29 Benguela Current	41 East-Central Australian Shelf	54 Chukchi Sea
6 Southeast U.S. Continental Shelf 7 Northeast U.S. Continental Shelf 20 Barents Sea 33 Red Sea 45 Northwest Australian Shelf 58 Kara Sea 8 Scotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland Sea 35 Gulf of Thailand 47 East China Sea 60 Faroe Pl 10 Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic 11 Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Sea	4 Gulf of California	17 North Brazil Shelf	30 Agulhas Current	42 Southeast Australian Shelf	55 Beaufort Sea
7 Northeast U.S. Continental Shelf 20 Barents Sea 33 Red Sea 45 Northwest Australian Shelf 58 Kara Sea 85 Scotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland Sea 9 Newfoundland-Labrador Shelf 22 North Sea 35 Gulf of Thailand 47 East China Sea 60 Faroe Pl 10 Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic 11 Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Sea	5 Gulf of Mexico	18 West Greenland Shelf	31 Somali Coastal Current	43 Southwest Australian Shelf	56 East Siberian Sea
8 Scotian Shelf 21 Norwegian Shelf 34 Bay of Bengal 46 New Zealand Shelf 59 Iceland S 9 Newfoundland-Labrador Shelf 22 North Sea 35 Gulf of Thailand 47 East China Sea 60 Faroe Pl 10 Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic 11 Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Se	6 Southeast U.S. Continental Shelf	19 East Greenland Shelf	32 Arabian Sea	44 West-Central Australian Shelf	57 Laptev Sea
9 Newfoundland-Labrador Shelf 22 North Sea 35 Gulf of Thailand 47 East China Sea 60 Faroe Pl 10 Insular Pacific-Hawaiian 23 Baltic Sea 36 South China Sea 48 Yellow Sea 61 Antarctic II Pacific Central-American Coastal 24 Celtic-Biscay Shelf 37 Sulu-Celebes Sea 49 Kuroshio Current 62 Black Se	7 Northeast U.S. Continental Shelf	20 Barents Sea	33 Red Sea	45 Northwest Australian Shelf	58 Kara Sea
10Insular Pacific-Hawaiian23Baltic Sea36South China Sea48Yellow Sea61Antarctic11Pacific Central-American Coastal24Celtic-Biscay Shelf37Sulu-Celebes Sea49Kuroshio Current62Black Se	8 Scotian Shelf	21 Norwegian Shelf	34 Bay of Bengal	46 New Zealand Shelf	59 Iceland Shelf
11 Pacific Central-American Coastal     24 Celtic-Biscay Shelf     37 Sulu-Celebes Sea     49 Kuroshio Current     62 Black Se	9 Newfoundland-Labrador Shelf	22 North Sea	35 Gulf of Thailand	47 East China Sea	60 Faroe Plateau
	0 Insular Pacific-Hawaiian	23 Baltic Sea	36 South China Sea	48 Yellow Sea	61 Antarctic
12 Caribbean Sea 25 Iberian Coastal 38 Indonesian Sea 50 Sea of Japan 63 Hudson	I Pacific Central-American Coastal	24 Celtic-Biscay Shelf	37 Sulu-Celebes Sea	49 Kuroshio Current	62 Black Sea
	2 Caribbean Sea	25 Iberian Coastal	38 Indonesian Sea	50 Sea of Japan	63 Hudson Bay
13 Humboldt Current 26 Mediterranean Sea 39 North Australian Shelf 51 Oyashio Current 64 Arctic O	3 Humboldt Current	26 Mediterranean Sea	39 North Australian Shelf	51 Oyashio Current	64 Arctic Ocean

## Wider Caribbean South-East Pacific South-West Atlantic LMEs 19,20,21,22,24,25,59,60.. North-East Atlantic






lobal Map of average primary productivity estimates based on SeaWiFS satellite data, with highs in red and lows in purple, and the boundaries of the 64 LMEs. (www.edc.uri.edu/lme)

## **UNEP Regional Seas Programme is Linked to** the LME 5-Module Suite of Ecosystem Condition Indicators to Support Management Actions

Large Marine Ecosystems (LMEs) are providing countries with a framework for progress in the direction of ecosystem-based adaptive management by recognizing the basic linkages between scientific assessments, protection of the marine environment, sustainable development of coastal and marine resources, and poverty alleviation.

The 5-module LME approach has developed indicators of productivity, fish and fisheries, pollution and ecosystem health, socioeconomics and governance to analyze ecosystem-wide changes. **Productivity** indicators measure the carrying capacity of an ecosystem for supporting living marine resources. The Fish and Fisheries module conducts assessments of dominant species within fish communities; and considers effects of naturally occurring environmental shifts in climate regime and excessive fishing effort causing shifts in species composition and abundance. Indicators in the Pollution and Ecosystem Health module help assess changes in coastal waters, estuaries and wetlands, and highlight eutrophic conditions. The **Socioeconomics** module examines how a sustainable marine resource base can meet the nutritional, social, economic and developmental needs of humans living in LME border countries. The **Governance** module engages multiple scales of national, regional and local jurisdictional frameworks needed to select and support ecosystem-based management practices leading to sustainable use of resources. The 5-modular approach, which provides the scientific foundation for management actions in ongoing GEF-LME Projects, can also be applied to the management of the 18 Regional Seas.


## **Modular Assessments for Sustainable Development**



### **Recent Trends in LMEs within Regional Seas,** identified through 5-Modular Assessments

### **Need for Precautionary Approach: Encroachment of Industrial Globalized Fisheries on Artisanal Fisheries in the Guinea Current LME**

K.A. Koranteng (2002) conducted a time series analysis of Catch-Per-Unit-Effort for both small-sized inshore artisanal-type vessels and industrialized fishing fleets from the European Union. The industrialized trawlers are fishing species in areas previously beyond technological and economic reach to support a globalized fishmeal extraction industry, utilized in industrialized farms in the developed world as animal feed or fertilizer. Koranteng found a consistent rise in industrial trawling and a downward trend in inshore seasonal artisanal fishing, which raises concerns for the fish harvest available to meet the growing nutritional needs of the 300 million people living along the Guinea Current coast.




Evolution of trawling effort in the Ghanaian demersal fisheries, by K.A. Koranteng, 2002.

### **Need for improved forecasts of fishery** fluctuations:

Toward long-term sustainability of pelagic and demersal fish stocks in the Humboldt Current LME


The Humboldt Current LME contains the world's largest upwelling system and is the world's most productive marine ecosystem, providing about 15-20% of the world's annual marine catch. Anchovy, sardine and horse mackerel are used for fish meal and for human consumption. Fishing sustains thousands of fishermen and their families. While the high productivity of the Humboldt Current LME is the result of upwelling processes governed by strong southerly trade winds, the upwelling is subjected to considerable interannual climatic variability, which causes variations in marine populations and catch. The normal seasonal upwelling can be interrupted by the El Niño-Southern Oscillation (ENSO) which results in intrusions of warm water. For the long-term sustainability of the pelagic and demersal fish stocks of this LME, improved forecasts of climate-driven fishery fluctuations are required.



### **Need to curb excessive nitrogen loading:** Model predictions of nitrogen affecting LMEs show significant increase

Excessive levels of nitrogen contributing to coastal eutrophication constitute a growing global environmental problem that is cross-sectoral in nature. Excessive nitrogen loadings have been identified as problems in the Baltic Sea, Black Sea, Adriatic portion of the Mediterranean, Yellow Sea, South China Sea, Bay of Bengal, Gulf of Mexico, and Patagonian Shelf LMEs. Model-predicted global estimates of dissolved inorganic nitrogen (DIN) export from freshwater basins to coastal waters in 1990 and 2050 have been developed by Kroeze and Seitzinger, 1999. These estimates, based on a business-as-usual (BAU) scenario, are cause for concern for future sustainability. Given the expected future increases in population and in fertilizer use, without significant nitrogen mitigation efforts, LMEs will be subjected to a future of increasing harmful algal bloom events, reduced fisheries, and hypoxia that will further degrade marine biomass and biological diversity.





Model-predicted nitrogen (dissolved inorganic N) export by rivers to coastal systems in 1990 and in 2050 — based on a business-as-usual (BAU) scenario. Figure modified from Kroeze and Seitzinger (1999).

# 我们的产品



## 大数据平台

国内宏观经济数据库

国际经济合作数据库

行业分析数据库

## 条约法规平台

国际条约数据库

国外法规数据库

## 即时信息平台

新闻媒体即时分析

社交媒体即时分析

## 预览已结束, 完整报告链接和二组

https://www.yunbaogao.cn/report/index/report?repor