


Carbon, biodiversity and land-use in the Central Congo Basin Peatlands

Lera Miles^a, Corinna Ravilious^a, Shaenandhoa García-Rangel^a, Xavier de Lamo^a, Greta Dargie^b and Simon Lewis^b

^a UN Environment World Conservation Monitoring Centre ^b, University of Leeds and University College London

KEY MESSAGES

- In January 2017, scientists announced new discoveries showing that the Central Congo Basin Peatlands form the largest area of peat swamp forest in the tropics. Estimates of soil organic carbon stocks in the Republic of Congo and Democratic Republic of Congo have radically increased as a result, to some 30 Gt. Degradation of these peat carbon stocks could have serious impacts on climate.
- Most of the area is covered by oil and gas concessions (for exploration and/or extraction), with smaller areas also covered by timber concessions and concessions for oil palm development or agricultural use. There is an associated risk of forest degradation, including from road development and woodfuel harvest.
- Concerns have been raised that the pattern of peatland drainage for oil palm plantations seen in Southeast Asia could also develop in the Central Congo Basin. While remote sensing provides good information about recent tree cover change in the area, less is understood about the potential scale of future land-use change and related risks.
- A Transboundary Ramsar Site was declared in June 2017, covering much of the Central Congo Basin Peatlands. The peat swamp forest ecosystem is in good condition due to its relative inaccessibility, and this new designation could provide a basis for its conservation and wise use consistent with the Ramsar Convention on wetlands.
- Further effective action to conserve and sustainably manage these peat swamp forests will also contribute to achieving the Sustainable Development Goals (as well as related national objectives):
 - Goal 13 on action to mitigate climate change, by keeping carbon stocks in the ground.
 - Goal 3 on health, by avoiding the serious air pollution problems observed in Southeast Asia when drained peatlands burn for months.
 - Goal 6 on clean water, through its target on protecting water-related ecosystems.
 - Goal 15 on life on land, by ensuring the conservation and wise use of this ecosystem and its threatened species.



The peatlands of the Congo Basin

A large, carbon-rich and relatively undisturbed peatland area within the central Congo Basin has caught the world's attention, following the publication of a key Nature paper in January 2017 [1]. These peat swamp forests are located in the heart of Africa and are shared between the Republic of Congo and the Democratic Republic of Congo (Map 1). They cover 40% of the Cuvette Centrale depression, occupying land between tributaries of the Congo River. This is now thought to be the largest tropical peatland area in the world. The discovery of their extent and significant carbon stocks is especially relevant to climate change mitigation [2]–[4], but also to the Sustainable Development Goals on health, water and life on land. As a unique ecosystem hosting both threatened species and endemic species only found in the Congo Basin region, its conservation would contribute to achieving the Aichi Biodiversity Targets (Map 2).

Peat is compressed, partly decayed organic matter under waterlogged conditions that prevent its full decomposition [5], [6]. Peatlands store about one-third of all soil carbon, despite covering only 3% of Earth's land area [7], [8]. They occur primarily within the tropics, the sub-arctic and boreal regions, and can host forests, grasslands, marshes and shrublands [3], [4]. Around 15% of global peatlands have already been disturbed (and/or drained) by agriculture, forestry, peat extraction and/or infrastructure development [4], [9].

The large-scale drainage of peatlands accounts for more than 3% of all greenhouse gas emissions [4]. When peatland are drained, the decomposition of organic matter resumes in peatlands releasing greenhouse gas emissions for decades or more (Figure 1) [9]. Drained tropical peatlands are also very vulnerable to fire, which can continue underground for months and contributes to haze with severe implications for air quality and human health, as well as to climate change [10]. The peatlands of the Central Congo Basin are waterlogged swamp forests, which have remained largely




Figure 1. Environmental and social impacts of peatland drainage © FAO 2017





Figure 2. Soil organic carbon stocks estimates in Central Congo Basin, Tropical and Global peatlands [1-3].

intact, with limited accessibility [1]. Although relatively undisturbed at present, these ecosystems may in future face similar threats to those reported now in Southeast Asia and across the tropics [11], [12]. Expansion of the road network over the past 15 years has increased access to previously remote locations (Map 3). Concessions for logging, mining and oil and gas exploration/exploitation together cover most of the region (Maps 3 and 4). As global food demand increases, agricultural expansion may also promote deforestation, peatland drainage and overall ecosystem degradation. Furthermore, some regional climate projections forecast reduced annual rainfall and stronger dry seasons, which could also facilitate peatland drainage [1].

Calls have already been made for the protection of the Central Congo Basin Peatlands whilst supporting national development needs [1]. The Republic of Congo recognized the role of peatland carbon stocks in the country's forest reference emission level for REDD+ [15]. Its Intended Nationally Determined Contribution under the climate change convention references a Permanent Forest Domain, in which land-use change would be illegal [29]. This includes a conservation forest category, in addition to the protected areas designated for biodiversity conservation [30]. This already represents a substantial commitment to protect the area.

The Republic of Congo is also considering the expansion of the *Lac Télé* Community Reserve to protect further areas of swamp forest (Map 1) [1]. The draft of its National REDD+ Strategy aims to ensure that agro-industrial concessions are not granted near wetlands or forests with high biodiversity value. REDD+ could be a potential tool to promote the conservation of the Congo peatlands. It was devised by the United Nations Framework Convention on Climate Change (UNFCCC), to support developing countries to reduce emissions from deforestation and forest degradation whilst ensuring the sustainable management of forest ecosystems and the conservation and enhancement of forest carbon stocks. The Convention's safeguards for REDD+ expect that its implementation delivers social and environmental benefits beyond climate change mitigation.

Map 1. Peatland extent, carbon content and protection status



According to the UK-Congolese team working on the Cuvette Centrale region, this is the most extensive tropical peatland complex worldwide, five times larger than previously estimated [1]. Initial estimates indicate an area of about 145,500 km², corresponding to 4% of the entire Congo Basin (Map 1) [1]. These estimates, however, require further ground-truthing as they rely on peat-vegetation associations and remote sensing products [1]. If validated, these findings could make the Democratic Republic of Congo the country with the second largest peatland extent (90,800 km²) in the tropics (after Indonesia), followed by the Republic of Congo (54,700 km²) [1]. The median peat depth is 2 metres (with a maximum of 5.9 metres), shallower than many other tropical peatlands, but with a higher carbon density in each handful of peat [1]. The Congo peat swamp forests store almost 30% of all the soil organic carbon found within tropical peatlands (about 30 billion tonnes of carbon), and about 5% of that estimated to be stored in peatlands across the world (Figure 2) [1]. Their carbon stocks are also estimated to be as high as those in the entire above- and below-ground forest biomass of the two countries [1] (Figure 3).

The two countries agreed in June 2017 to cooperate in the sustainable management of three Ramsar sites that together occupy 45% of the peatland area, by formally declaring them as a single transboundary site, Complexe Transfrontalier lac Télé - Grands Affluents - Lac Tumba. Only 7% of the peatland region is covered by designated protected areas at the national level, so this transboundary site provides a good framework for conservation and wise use of the area consistent with the Ramsar Convention on wetlands.




Figure 3. Carbon stock estimates for peat and above- and below-ground woody biomass across the Republic of Congo and the Democratic Republic of Congo [1], [30].

Map 1 data sources:

Peat swamp forest vegetation: Dargie GC, Lewis SL, Lawson IT, Mitchard ETA, Page SE, Bocko YE, Ifo SA. 2017. Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature* 542:86–90. Map data available from www.afritrn.org/en/peatland.

Protected areas: IUCN and UNEP-WCMC. 2017. The World Database on Protected Areas (WDPA). Public release July 2017. Cambridge, UK. Available at: www.protectedplanet.net. Please note: some Ramsar site boundaries have been digitized by UNEP-WCMC.


Map 2. Richness of threatened vertebrates within the Central Congo Basin Peatlands



The Central Congo Basin Peatlands are home to 14 species threatened at the global level, as well as to 10 prioritised nationally and/or regionally [20], [21]. For example, these include three of the four African ape species. To the east of the Congo river, bonobos (*Pan paniscus*) live in the peat swamp forests, while to the west, western lowland gorillas (*Gorilla gorilla gorilla*) can be found [22]. Chimpanzees (*Pan troglodytes*) are found in the north of the Central Congo Basin peatlands [22]. Swamp forest provides them with useful food resources, with both gorillas and bonobos enjoying aquatic herbs. Map 2 shows the known richness of threatened vertebrates across the Congo peatlands. A total of 5 areas have been identified as key for biodiversity conservation within this region (Map 2).

All the African apes are protected by the Kinshasa Declaration on Great Apes as well as under national law [23]. At a regional scale,

the COMIFAC Convergence Plan, in its “Axis of intervention for the conservation and enhancement of biodiversity” emphasizes the importance of the conservation of large mammals and other threatened species of fauna and flora [24].




Map 2 data sources:


Peat swamp forest vegetation: Dargie GC, Lewis SL, Lawson IT, Mitchard ETA, Page SE, Bocko YE, Ifo SA. 2017. Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature* 542:86–90. Map data available from www.afritrn.org/en/peatland.

Threatened vertebrate species richness: IUCN 2015. The IUCN Red List of Threatened Species. Version 2015.1. <http://www.iucnredlist.org>. Downloaded on 23/2/17.

Map 3. Forest concessions across the Central Congo Basin peatlands, showing tree cover loss along logging roads and elsewhere




Most of the Central Congo Basin Peatlands are relatively undisturbed, but over the past ten years, tree cover has been lost at the edge of the peatlands and along logging roads (Map 3). This has been mainly within the Democratic Republic of Congo, where reports indicate that slash-and-burn agriculture has been the main driver of this change [16]–[18]. A considerable number of roads have also been opened within this period (Map 3). This may facilitate access to previously remote peat swamp forests. About 28,956 km² (20%) of peatlands are within forest concessions, and approximately 53% of these are thought to be already allocated to operators [25].



Map 3 data sources:

Peat swamp forest and other vegetation: Dargie GC, Lewis SL, Lawson IT, Mitchard ETA, Page SE, Bocko YE, Ifo SA. 2017. Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature* 542:86–90. Map data available from www.afritrn.org/en/peatland. **Tree cover loss:** Hansen, MC., Potapov, PV., Moore, R., Hancher, M., Turubanova, SA., Tyukavina, A., Thau, D., Stehman, SV., Goetz, SJ., Loveland, TR., Kommareddy, A., Egorov, A., Chini, L., Justice, CO, and Townshend, JRG. 2013. High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* 342 (15 November): 850–53. Data available on-line from: <http://earthenginepartners.appspot.com/science-2013-global-forest>. **Roads:** WRI. 2009. Interactive Forest Atlas Version 1.0 for Congo; DIAF, DGF and WRI. 2013. Interactive Forest Atlas Version 1.0 for Democratic Republic of Congo. **Logging Roads:** OpenStreetMap with contributions from the Logging Road Initiative. n.d. Congo Basin logging roads. Accessed through Global Forest Watch on 24/02/17. www.globalforestwatch.org. **Forest concessions:** World Resources Institute. n.d. Logging. Accessed through Global Forest Watch on 16/11/17. www.globalforestwatch.org. **Agricultural concessions:** DIAF/WRI. 2017. Agricultural concessions. Created based on the FACET data, 'cartes territoriales' and a repertory of plantations. Layer in the process of being validated. Accessed through Moabi DRC on <https://mapforenvironment.org/layer/info/80/DRC-Agriculture-Concessions#5.67/-1.263/19.467> on 03/03/17.

Map 4. Mining permits and oil and gas concessions in peat swamp forest



Oil and gas concessions cover the Central Congo Basin Peatlands almost in its entirety (Map 4). Licenses have already been granted across 30% of the area (Republic of Congo 31,288 km², Democratic Republic of Congo 12,990 km²), while 13% appears to be still open for bidding [26], [27]. The status of 44% of the oil and gas concessions in the Democratic Republic of Congo was unknown

as peat has a high greenhouse gas emission intensity and slow renewal rates [31].



预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_14488

