



Food and Agriculture
Organization of the
United Nations



International Fund for
Agricultural Development

unicef



World Food
Programme



World Health
Organization

2020

THE STATE OF FOOD SECURITY AND NUTRITION IN THE WORLD

TRANSFORMING FOOD SYSTEMS
FOR AFFORDABLE HEALTHY DIETS

This flagship publication is part of **THE STATE OF THE WORLD** series of the Food and Agriculture Organization of the United Nations.

Required citation:

FAO, IFAD, UNICEF, WFP and WHO. 2020. *The State of Food Security and Nutrition in the World 2020. Transforming food systems for affordable healthy diets*. Rome, FAO.
<https://doi.org/10.4060/ca9692en>

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO), the International Fund for Agricultural Development (IFAD), the United Nations Children's Fund (UNICEF), the World Food Programme (WFP) or the World Health Organization (WHO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO, IFAD, UNICEF, WFP or WHO in preference to others of a similar nature that are not mentioned.

The designations employed and the presentation of material in the maps do not imply the expression of any opinion whatsoever on the part of FAO, IFAD, UNICEF, WFP or WHO concerning the legal or constitutional status of any country, territory or sea area, or concerning the delimitation of frontiers.

All reasonable precautions have been taken by FAO, IFAD, UNICEF, WFP and WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall FAO, IFAD, UNICEF, WFP and WHO be liable for damages arising from its use.

ISSN 2663-8061 (PRINT)
ISSN 2663-807X (ONLINE)
ISBN 978-92-5-132901-6
© FAO 2020



Some rights reserved. This work is made available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, this work may be copied, redistributed and adapted for non-commercial purposes, provided that the work is appropriately cited. In any use of this work, there should be no suggestion that FAO endorses any specific organization, products or services. The use of the FAO logo is not permitted. If the work is adapted, then it must be licensed under the same or equivalent Creative Commons licence. If a translation of this work is created, it must include the following disclaimer along with the required citation: "This translation was not created by the Food and Agriculture Organization of the United Nations (FAO). FAO is not responsible for the content or accuracy of this translation. The original English edition shall be the authoritative edition."

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the Arbitration Rules of the United Nations Commission on International Trade Law (UNCITRAL) as at present in force.

Third-party materials. Users wishing to reuse material from this work that is attributed to a third party, such as tables, figures or images, are responsible for determining whether permission is needed for that reuse and for obtaining permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

Sales, rights and licensing. FAO information products are available on the FAO website (www.fao.org/publications) and can be purchased through publications-sales@fao.org. Requests for commercial use should be submitted via: www.fao.org/contactus/licence-request. Queries regarding rights and licensing should be submitted to: copyright@fao.org.

COVER PHOTOGRAPH ©Mint Images

THAILAND. Floating market on a canal in Bangkok, where local boats are laden with fresh food.

2020

**THE STATE OF
FOOD SECURITY
AND NUTRITION
IN THE WORLD**

**TRANSFORMING FOOD SYSTEMS
FOR AFFORDABLE HEALTHY DIETS**



Food and Agriculture Organization of the United Nations
Rome, 2020

CONTENTS

FOREWORD	viii	ANNEXES	163
METHODOLOGY	xi	ANNEX 1A	
ACKNOWLEDGEMENTS	xii	Statistical tables to Part 1	164
ACRONYMS AND ABBREVIATIONS	xiv	ANNEX 1B	
KEY MESSAGES	xvi	Methodological notes for the food security and nutrition indicators	190
EXECUTIVE SUMMARY	xviii	ANNEX 2	
		Methodologies Part 1	202
PART 1		ANNEX 3	
FOOD SECURITY AND NUTRITION AROUND THE WORLD IN 2020	1	Description, data and methodology of Section 2.1	214
1.1 Progress towards hunger and food insecurity targets	3	ANNEX 4	
1.2 Progress towards global nutrition targets	26	National food-based dietary guidelines (FBDGs) used to compute the cost of a healthy diet	231
1.3 The critical link between food security and nutrition outcomes: food consumption and diet quality	40	ANNEX 5	
1.4 Conclusions	60	Additional tables and figures to Section 2.1	236
PART 2		ANNEX 6	
TRANSFORMING FOOD SYSTEMS TO DELIVER AFFORDABLE HEALTHY DIETS FOR ALL	63	Definition of country groups	241
2.1 The cost and affordability of healthy diets around the world	65	ANNEX 7	
2.2 The hidden health and environmental costs of what we eat	93	Description, data and methodology of Section 2.2	242
2.3 What is driving the cost of nutritious foods?	115	ANNEX 8	
2.4 Policies to reduce the cost of nutritious foods and ensure affordability of healthy diets	138	Health and climate-change costs related to dietary risks	248
		ANNEX 9	
		Glossary	253
		NOTES	258

TABLES, FIGURES AND BOXES

TABLES

1 Prevalence of undernourishment (PoU) in the world, 2005–2019	9	9 Shifting to healthy diets that include sustainability considerations can contribute to reductions in environmental impacts on land, energy and water use	105	A2.1 Monitoring rules and classification of progress towards achieving the nutrition targets at the regional and global levels used in this report	204
2 Number of undernourished people in the world, 2005–2019	11	10 Fruits and vegetables show the highest levels of food price seasonality in seven selected countries in Africa (2000–2012)	123	A2.2 Data sources used to assess diets: strengths and limitations for global assessment	208
3 Prevalence of food insecurity at severe level only, and moderate or severe level, measured with the Food Insecurity Experience Scale, 2014–2019	20	11 Agricultural production in low-income countries is penalized, while in middle- and high-income countries it is supported (2005–2016)	132	A2.3 Food group classifications used in the analysis	210
4 Number of people experiencing food insecurity at severe level only, and moderate or severe level, measured with the Food Insecurity Experience Scale, 2014–2019	21	12 Worldwide, governments support the production of sugar, rice and animal products the most through interventions, while penalizing the production of more nutrient-rich fruits and vegetables (2005–2016)	133	A3.1 Nutrient intake values for a representative 30-year-old woman	216
5 The global nutrition targets endorsed by the World Health Assembly and their extension to 2030	26	A1.1 Progress towards the Sustainable Development Goals (SDGs) and global nutrition targets: Prevalence of undernourishment, moderate or severe food insecurity, selected forms of malnutrition, exclusive breastfeeding and low birthweight	164	A3.2 Cost and affordability of the three diets by country (170 countries), by region, country income group and population (millions) in 2017	223
6 Most regions are making some progress, but not enough to achieve global targets; no subregion is on track for the low birthweight target, and adult obesity is worsening in all subregions	30	A1.2 Progress towards the Sustainable Development Goals (SDGs) and global nutrition targets: Number of people who are affected by undernourishment, moderate or severe food insecurity and selected forms of malnutrition; number of infants exclusively breastfed and number of babies born with low birthweight	177	A3.3 Lower- and upper-bound estimates of the percentage and number of people (in millions) who cannot afford the cost of each diet, by region and country income group in 2017	229
7 The cost of a healthy diet is 60 percent higher than the cost of the nutrient adequate diet, and almost 5 times the cost of the energy sufficient diet in 2017	77			A4.1 Food-based dietary guidelines used to compute the cost of a healthy diet	233
8 More than 3 billion people in the world cannot afford a healthy diet in 2017	85			A5.1 Average estimates of own-price elasticities by food group	236
				A5.2 Predicted own- and cross-price elasticities by food group and country income group in 2008	237

TABLES, FIGURES AND BOXES

A7.1 Summary description of baseline current food consumption (BMK) and four alternative healthy diet patterns that include sustainability considerations (FLX, PSC, VGT and VGN)

243

A7.2 Overview of per capita food consumption (g/day) and kilocalorie intake (kcal/day) by diet scenario in year 2010 – global

245

A8.1 Percent contribution of diet- and weight-related risk factors to the reduction in mortality in year 2030 when moving from the benchmark diet to the four alternative healthy diet patterns – global

249

FIGURES

1 The number of undernourished people in the world continued to increase in 2019. If recent trends are not reversed, the SDG 2.1 Zero Hunger target will not be met

4

2 PoU in Africa by subregion, with projections to 2030. The highest levels of undernourishment are found in Middle and Eastern Africa

10

3 PoU in Asia by subregion, with projections to 2030. Western Asia has been off track in the recent past and is the only subregion in the continent where the prevalence of undernourishment is projected to increase

12

4 PoU in Latin America and the Caribbean by subregion, with projections to 2030. Projections point to convergence between Central America, where undernourishment is projected to increase, and the Caribbean, where it is projected to decline

5 If recent trends persist, the distribution of hunger in the world will change substantially, making Africa the region with the highest number of undernourished in 2030

6 Explanation of food-insecurity severity levels measured by the FIES in SDG Indicator 2.1.2

7 Moderate or severe food insecurity affects one quarter of the world population, and it has been increasing over the past six years. Over half of the population in Africa, almost one-third in Latin America and the Caribbean and more than one-fifth in Asia are food insecure

8 Over half of the people affected by moderate or severe food insecurity in the world live in Asia and more than one-third live in Africa

9 Globally and in every region, the prevalence of food insecurity is slightly higher in women than in men

10 Despite some progress for most indicators, only the 2025 target for exclusive breastfeeding is on track to be achieved. Childhood overweight and adult obesity trends need to be reversed

28

15

11 Most regions are not on track to achieve the targets for child overweight, and adult obesity is on the rise in all regions

32

16

12 Sub-Saharan Africa is the only subregion with a rising number of stunted children

35

19

13 Globally, 21.3 percent of children under 5 are stunted, and 7 subregions out of 17 have high or very high stunting prevalences in 2019

36

14 The prevalence of stunting is higher in rural areas and in poorest households

37

22

15 In Latin America and the Caribbean, stunting prevalence of children living in the poorest households is about three times higher compared to those living in the richest households

38

23

16 Different ways of applying the principles for healthy diets: examples from three countries

43

25

17 From 2000 to 2017, the global availability of fruits and vegetables and dairy products increased, while availability of staple foods remained stable

47

18	From 2000 to 2017, the global availability of pulses and nuts, and poultry and fish increased, while availability of red meat increased largely in upper-middle-income countries	48	25	The unaffordability of a healthy diet is strongly associated with food insecurity and different forms of malnutrition, including child stunting and adult obesity	68	32	Compared with current consumption patterns, adopting any of the four alternative healthy diet patterns is projected to lead to significant reductions in mortality by 2030	100
19	From 2000 to 2017, the global availability of processed meat, eggs and fats increased, while availability of sugar decreased	49	26	Income elasticities of demand for food products tend to be higher in countries where income per capita is lower	71	33	Under current food consumption patterns, diet-related health costs are projected to reach USD 1.3 trillion per year in 2030	101
20	The proportions of different food groups available for human consumption differ across country income groups: a snapshot of 2017	51	27	Dairy, fruits, vegetables and protein-rich foods are the highest-cost food groups for a healthy diet globally in 2017	79	34	Adoption of any of the four alternative healthy diet patterns would dramatically decrease diet-related health costs by 2030	102
21	Children living in urban households and those from richer families have better dietary diversity	54	28	A healthy diet is unaffordable for the poor in every region of the world in 2017	81	35	Adoption of any of the four alternative healthy diet patterns could significantly reduce projected diet-related GHG emissions in 2030	106
22	Percentage of children 6–23 months of age eating foods from minimum number of food groups. The vast majority of children 6–23 months of age are not meeting the minimum dietary diversity	55	29	In most countries in the Global South, the cost of a healthy diet exceeds average national food expenditures per capita in 2017	83	36	Increases in GHG emissions will be necessary for some countries to meet dietary energy and protein needs	108
23	The majority of children in the world consume grains, roots and tubers. Few children are being fed flesh foods or eggs	56	30	Affordability of a nutrient adequate diet varies widely within many countries due to temporal and geographical variations in prices and differences in incomes	89	37	Adoption of plant-based dietary patterns would reduce the social cost of GHG emissions by 41–74 percent in 2030	109
24	As food insecurity becomes more severe, food consumption and dietary energy intake decrease and staples make up a larger share of the diet	59	31	Pregnant and lactating women and adolescent girls have higher dietary energy and iron needs, which increases the cost of a nutrient adequate diet: Burundi and Uganda case studies	91	38	The adoption of any of the four alternative healthy diet patterns could potentially lead to 22–29 percent reduction in the full cost of diets on average by 2030	111

TABLES, FIGURES AND BOXES

39 Protectionary trade policies protect and incentivize domestic production of staple foods, such as rice, but often to the detriment of nutritious foods in low-income countries	134	A8.1 If current food consumption patterns continue, diet-related health costs are projected to be USD 1.3 trillion in 2030 – by country income group and cost component	249	6 How do current consumption patterns compare with national food-based dietary guidelines recommendations? Case study from Belgium	45
40 Public expenditure in food and agriculture favours producer subsidies, with less investment towards efficiency gains across food supply chains (selected African countries, 2005–2017)	136	A8.2 Adoption of any of the four alternative healthy diets could significantly reduce projected diet-related GHG emissions in 2030	251	7 Potential impacts of COVID-19 pandemic on availability and access to nutritious foods and overall diet quality	52
41 Policy options to reduce the cost of nutritious foods and enhance affordability of healthy diets with complementary policies to promote healthy diets	144	A8.3 The social cost of GHG emissions under current food consumption patterns for different emissions-stabilization scenarios in 2030 (USD billion)	251	8 Minimum dietary diversity for women: evidence from three countries	53
A4.1 Average cost of a healthy diet by applying ten different FBDGs definitions across all countries globally, and four different EAT-Lancet reference diet patterns in 2017	232	A8.4 Three-quarters of the social cost of GHG emissions under current food consumption patterns comes from meat and dairy products in 2030	252	9 Association between food insecurity based on the FIES and diet quality based on new metrics: evidence from Ghana and the United Republic of Tanzania	57
A5.1 Average percentage				10 Description of the three diets used in the cost and affordability analysis	73
		BOXES		11 Computing the lowest cost of the three diets used in the cost	

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_3396

