

TOBACCO & POSTSURGICAL OUTCOMES

- Those who smoke, or even quit smoking, before surgery have a significantly increased risk of experiencing a range of postsurgical complications compared with non-smokers.
- Children exposed to second-hand tobacco smoke have a higher risk of peri-anaesthetic respiratory adverse events.
- Those who quit smoking approximately 4 weeks before surgery have a reduced risk of postsurgical complications. The optimum length of cessation varies depending on the type of postsurgical outcome assessed (i.e. wound healing or total complications).
- High-intensity behavioural interventions, which include weekly contact, provision of nicotine or non-nicotine (varenicline) replacement therapy and referrals to telephone cessation support and which are delivered for at least four weeks before surgery, are effective in reducing postsurgical complications. The impact of such interventions on smoking cessation can occur within one-week pre surgery.
- The surgical unit can play a vital role in the assessment of smoking status and initiation of smoking cessation interventions in its patients before surgery.
- Governments should promote the implementation of smoke-free hospital policies, access to nicotine or non-nicotine (varenicline) replacement therapy and access to community tobacco cessation services for presurgical patients.

Tobacco use

Worldwide, more than 1.1 billion people smoke tobacco and at least 367 million people use smokeless tobacco (1). Tobacco smoking is known to cause adverse health effects, including cancer, cardiovascular and respiratory diseases (2), and was responsible for approximately 11.5% of total deaths in 2015 (3).

Surgery burden

Approximately one in 25 individuals (representing between 187 million and 280 million cases globally) undergoes major surgery annually for the treatment of disease, injury or illness (4, 5). Complications from surgery such as surgical site infections and respiratory and cardiopulmonary events represent a substantial burden for both patients and health-care systems. Major morbidity occurs in between 4% and 16% of all inpatient surgical procedures in developed countries, with perioperative mortality and severe disability occurring in 1% of cases (6-9). In developing countries, mortality rates reportedly increase to up to 0.24–10% of cases (9-11). Surgical procedures with major complications cost significantly more than surgery without any complications (12), suggesting that substantial savings can be made if such complications are prevented. Surgery and anaesthesia cause severe stress and trauma to the body (13).

Definitions

Smoked tobacco: any product made entirely or partly of leaf tobacco that is intended to be lit and the produced smoke inhaled. Examples include manufactured cigarettes, roll-your-own cigarettes, water pipes (e.g. hookah, shisha), cigars, kreteks and bidis.

Second-hand smoke (SHS): the combination of “mainstream” smoke (the smoke emerging from the mouth end of a cigarette during smoking) that is exhaled by the smoker, and “side-stream” smoke emitted into the environment from lit cigarettes and other tobacco products. The terms “Passive smoking or Involuntary smoking or Environmental tobacco smoke” are also often used to describe exposure to SHS.

Smokeless tobacco: any product that consists of cut, ground, powdered or leaf tobacco that is intended to be placed, loose or in sachets, in the oral or nasal cavity. Examples include snuff, chewing tobacco, gutka and mishri.

In the postsurgical period, the body undergoes a post-traumatic inflammatory response to fight infections and activates a wound-healing cascade for tissue recovery. The recovery process increases the body's need for oxygen and other nutrients, and modifiable risk factors including high body mass index, risky alcohol consumption and active smoking are thought to interfere with this process (14, 15).



This evidence brief aims (a) to summarize the association between tobacco exposure (smoking, smokeless and second-hand smoke) and postsurgical complications; and (b) to describe the effectiveness of interventions to reduce presurgical tobacco use and tobacco-related complications.

The impact of tobacco use on postsurgical outcomes

Chronic exposure to tobacco causes adverse physiological changes in cardiovascular function, pulmonary function and tissue healing. These changes may interfere with the postsurgical recovery process and account for the increased occurrence of postsurgical complications observed in smokers (13). Additionally, there is some evidence that smoking even one cigarette can result in reduced blood flow which, in turn, can contribute to adverse surgical outcomes (16, 17).

Cardiovascular function Chemical substances contained in tobacco increase the body's need for oxygen, but reduce its capacity to use oxygen (18). Nicotine stimulates the central nervous system, increasing blood pressure, heart rate, peripheral vascular resistance and oxygen consumption. Nicotine is also thought to induce vasoconstriction and inhibit platelet aggregation, reducing oxygen transport. Carbon monoxide reduces the availability of oxygen for cellular processes by binding to haemoglobin, and also inactivates cardiac enzymes, leading to decreased oxygen transport and use (19). Together, these result in tissue hypoxia and increased blood viscosity, which increases an individual's risk of cardiovascular events.

Pulmonary function Smoking has an adverse impact on pulmonary function, primarily through decreased mucociliary clearance and abnormal small airway function (20). Cigarette smoking damages the ciliated epithelium and the tracheobronchial tree in the lungs, leading to increased mucus, obstruction in the bronchioles and reduced ciliary function, and increases the risk of infections and respiratory complications (14). Mucous hypersecretion leads to increased sputum volume, which may result in deterioration in the oxygen transport system, inflammation of the airway and increased pulmonary complications.

Impaired wound healing Smoking may impair surgical site healing and promote wounds opening along sutures (dehiscence) via a number of pathways including (a) peripheral tissue hypoxia leading to necrosis; (b) decreased inflammatory responses; and (c) delayed proliferative healing responses and reduced collagen synthesis (14). Increased oxidative stress inhibits the mechanisms of neutrophils, which slows down the wound-healing process and reduces the body's capacity to fight bacterial infections (21). Smoking also impairs production of pro- and anti-inflammatory cytokines responsible for regulating the immune function within the body, which may be a predisposing risk factor for infections in the postoperative period (14).

Impaired bone healing Smoking may also affect bone healing in several ways, including increased tissue hypoxia, vasoconstriction secondary to nicotine and direct impairment of osteoblast activity and collagen synthesis by tobacco smoke (22), with a systematic review identifying smoking as one of the top 10 risk factors for non-union of long bones (23).

Second-hand smoke exposure and smokeless tobacco Little is known about the ways smokeless tobacco may influence postsurgical outcomes. In children, environmental exposure to tobacco smoke is associated with significantly higher odds of adverse surgical outcomes.

Association between tobacco exposure and postsurgical complications – reviews of observational trials

The link between tobacco smoking and the presence of postsurgical complications has been well studied. Overall, 28 systematic reviews published since 2004 were identified, examining the impact of smoking on a range of postsurgical outcomes (see Annex 1). All reviews reported that smoking, even when the smoker quits before surgery, was significantly associated with increased risk of at least one adverse postsurgical outcome, compared with the results for non-smokers (see Annex 1). A 2014 review (26) of 107 observational studies found a positive association between preoperative smoking status and

TOBACCO & POSTSURGICAL OUTCOMES

a number of postoperative complications (within 30 days of surgery). The confounder-adjusted relative risks of surgical/intraoperative or postoperative complications were significantly higher in smokers in the case of: general morbidity/total complications (RR: 1.75, 95% CI: 1.40–2.20), wound complications (RR: 2.49, 95% CI: 1.91–3.26), general infections (RR: 2.05, 95% CI: 1.34–3.13), pulmonary complications (RR: 2.46, 95% CI: 1.74–3.48); neurological complications (RR: 1.71, 95% CI: 1.07–2.74); and admission to an intensive care unit after surgery (RR: 1.6, 95% CI: 1.14–2.25) (26). For wound healing in particular, a review of 140 cohort studies involving 479 150 patients found that smokers had significantly higher adjusted odds ratios for healing delay (OR: 2.07, 95% CI: 1.53–2.81) and dehiscence (OR: 1.79, 95% CI: 1.57–2.04), surgical site infection (OR: 2.27, 95% CI: 1.82–2.84) wound complications in hernia (OR: 2.07, 95% CI: 1.23–3.47), and lack of fistula or bone healing (OR: 2.44, 95% CI: 1.66–3.58) compared with non-smokers. These findings of increased risk of experiencing postsurgical complications are consistent with other reviews which include patients across all surgical specialities (14, 27–30), as well as reviews which include only patients undergoing surgery in specific sites, including hip and knee (31, 32), operation for Crohn's disease (33), lower extremity grafting (34), periodontal surgery (35), spinal surgery (15), inguinal hernia surgery (36) and hip arthroplasty (37). The risk of delayed wound healing is similarly elevated in cosmetic surgery (OR: 2.50, 95% CI: 0.49–4.08) and bariatric surgery patients (OR: 3.30, 95% CI: 1.90–5.64) (38).

Association between cessation and postsurgical complications – reviews of controlled trials

A review of randomized controlled trials showed that interventions to increase cessation can significantly reduce the incidence of any postsurgical complication (RR: 0.42, 95% CI: 0.27–0.65) (39) and surgical site infections (OR: 0.43, 95% CI: 0.21–0.85) (14), and postoperative morbidity up to six months post-follow-up (40).

Association between exposure to second-hand smoke or use of smokeless tobacco and postsurgical complications

A systematic review and meta-analyses examining the impact of environmental tobacco smoke exposure on anaesthetic and surgical outcomes in children found that exposure significantly increased risk of peri-anaesthetic respiratory adverse events (RR: 2.52, 95% CI: 1.68–3.77) (41). Observational studies suggest that second-hand smoke exposure may be associated with adverse respiratory outcomes in both adults and children during general anaesthesia, as well as prolonged recovery time (42–47).

Association between tobacco exposure and postsurgical complications – summary of findings

Evidence from systematic reviews of observational studies shows a significantly increased risk of postoperative complications in smokers for all types of surgery, as well as in specific surgical sites including hip and knee, bowel resection and spinal surgery (40). The association between quitting smoking approximately 3–4 weeks before surgery and reduced postoperative complications has also been consistently reported in systematic reviews of randomized controlled trials (21).

Are there increased risks associated with short-term smoking cessation prior to surgery?

While there is a general consensus that stopping smoking before surgery can improve outcomes, there has been some controversy about the optimal timing of smoking cessation. Two studies published by Warner and colleagues (48–50) in a small sample of patients (< 200) undergoing coronary artery bypass grafting reported higher, but non-significant, rates of pulmonary complications among those who stopped smoking less than eight weeks prior to surgery, compared with those who continued to smoke.



Evidence from systematic reviews of observational studies has, however, reported no increase in adverse outcomes in people who cease smoking less than eight weeks before surgery (all complications: RR: 0.78, 95% CI: 0.57–1.07) (29), between two and four weeks (pulmonary complications: RR: 1.14, 95% CI: 0.90–1.45) and less than two weeks (pulmonary complications: RR: 1.20, 95% CI: 0.96–1.50) presurgery, compared with current smokers (27, 28). Longer abstinence periods (> 4 weeks) are, however, consistently associated with better postsurgical outcomes (21, 27, 28, 51), with a review reporting that each additional week of cessation resulted in an improvement of 19% in terms of reduction of postoperative morbidity (28).

Effectiveness of interventions to reduce presurgical tobacco use and related complications

Given the benefits of cessation for the reduction of adverse outcomes related to surgery, the presurgical period represents a key opportunity for interventions to reduce smoking and related complications. In our search, nine systematic reviews examining interventions to reduce smoking in patients undergoing surgery were identified. This includes a Cochrane systematic review published in 2014, which identified 13 randomized controlled trials examining interventions to reduce smoking in the presurgical period. Both low-intensity (RR: 1.30, 95% CI: 1.16–1.46) and high-intensity (RR: 10.76, 95% CI: 4.55–25.46) behavioural interventions were effective in reducing smoking immediately following the intervention. Only the two high-intensity interventions, which included weekly contacts (52, 53), were effective in reducing smoking rates at 12 months' follow-up (RR: 2.96, 95% CI: 1.57–5.55), with a corresponding reduction in postoperative complications (RR: 0.42, 95% CI: 0.27–0.65) (39). A narrative review of the efficacy of nicotine replacement therapy (NRT) in the perioperative period found limited evidence to indicate any increased risk of healing-related or cardiovascular-related complications (54). Only one trial examined the impact of non-NRT pharmacological interventions (varenicline), and found an increase in cessation

rates at 12 months' follow-up (RR: 1.45, 95% CI: 1.01–2.07), but no reduction in risk of postsurgical complications (RR: 0.94, 95% CI: 0.52–1.72) (55). Since the publication of the Cochrane review, other randomized controlled trials have found that a lower-intensity intervention that did not require weekly face-to-face sessions (NRT plus a telephone quitline), were effective in reducing smoking rates at 12 months' follow-up (RR: 3.0, 95% CI: 1.2–7.8) (56). Further, in a randomized controlled trial with non-NRT pharmacological interventions (varenicline) and telephone quitline follow-up, compared with a brief intervention without pharmacotherapy, an increase in cessation rates was found at 12 months' follow-up (1.62, 95% CI: 1.16–2.25, $P = 0.003$) (57). Consistent with reviews in hospitalized (58) and non-hospitalized populations (59), a small number of studies suggest that high-intensity behavioural interventions, with weekly face-to-face or telephone contact, the offer of NRT and referrals to the quitline, provided at least four weeks before surgery, are effective in reducing smoking and postoperative complications in presurgical patients.

Potential next steps

Research

There is strong evidence indicating that tobacco smoking is associated with an increased risk of a range of adverse postoperative outcomes. However, more prospective research, with longer follow-up times, is needed to assess the impact of smokeless tobacco use on these outcomes. Because of the negative impact of nicotine on cardiovascular function, research is also needed on the potential impact of use of electronic nicotine delivery systems on surgical outcomes, since these products can contain high and variable levels of nicotine. Finally, although current evidence suggests that effective intervention strategies exist for the reduction of smoking in preoperative patients, these results are based on a small number of trials conducted in developed countries. There has been only a small number of trials examining the use of pharmacotherapy (non-NRT) on smoking cessation in these patients. More intervention research examining different intervention intensities and

TOBACCO & POSTSURGICAL OUTCOMES

modalities and the usefulness of pharmacotherapy is needed to inform future strategies to reduce smoking in this population.

Practice

Although a relatively large number of patients would like to quit smoking, and scheduled surgery provides a potential teaching opportunity to help smokers quit in the long term, patients are often poorly informed of the benefits of smoking cessation for surgical outcomes and unaware of the resources available to help them to quit (60). There is potential for surgeons and anaesthetists to be involved in the initiation and delivery of preoperative smoking cessation care. While their capacity to offer high-intensity behavioural interventions is likely to be limited, surgical staff can play an active role by identifying smokers and assessing their willingness to quit smoking, providing information on the potential implications of continuing to smoke for surgical outcomes, supporting the initiation of NRT and referring patients to cessation services.

Systems

Perioperative services have a considerable role to play in supporting smoking cessation among surgical patients (61). However, the involvement of other hospital staff, primary care physicians and community resources may help to support such services by ensuring that a comprehensive and individualized smoking cessation intervention is developed before a planned operation (60, 62). The introduction of smoke-free policies in hospitals such as those encouraged by the WHO Framework Convention on Tobacco Control is essential to facilitate efforts to reduce smoking in presurgical patients. Such policies should be accompanied by the provision of inpatient cessation care (in the form of brief advice and provision of NRT in preoperative and postoperative units) as well as outpatient community cessation care (in the form of tobacco quitlines and active referrals to primary care physicians for advice and/or NRT or non-NRT pharmacotherapy). Mechanisms to facilitate active referral to such cessation resources should be implemented as part of the provision of routine surgical care in order to ensure the best outcomes for surgical patients.

Methods

WHO conducted a systematic search of the peer-reviewed literature for systematic reviews that examined (a) the impact of reducing the use of smoked or smokeless tobacco and exposure to second-hand smoke on postsurgical outcomes; and (b) interventions to reduce preoperative smoking, published between 2004 and May 2016. To make the review as inclusive as possible, a broad definition of systematic review was used. Specifically, systematic reviews were defined as reviews that included defined inclusion/exclusion criteria and provided information to indicate that a systematic method of searching and selection of trials had been undertaken. Where insufficient evidence from systematic reviews was present, a selective examination of non-systematic reviews and high-quality studies were undertaken. Relevant high-quality systematic reviews known to the authors and undertaken outside the search period were also included. The inclusion criteria included humans only; subjects exposed to any form of tobacco who were scheduled to undertake surgery; and examination of association with and/or reduction of smoking in relation to postsurgical outcomes. The review was not limited by language, although few non-English studies were identified. Using standardized methods, one reviewer undertook data extraction for information pertinent to the objectives of the evidence brief. Findings from any meta-analyses (relative risk (RR)/odds ratios (OR) and corresponding 95% confidence interval (CI)) undertaken within the relevant reviews were reported within the text or in a supplementary table, where available.

Tobacco and postsurgical outcomes: WHO tobacco knowledge summaries/ Serene L Yoong, Edouard Tursan d’Espaignet, John Wiggers, Simone St Claire, Jannicke Mellin-Olsen, Alice Grady et al.

ISBN 978-92-4-000036-0 (electronic version)

ISBN 978-92-4-000972-1 (print version)

© World Health Organization 2020

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO licence (CC BY-NC-SA 3.0 IGO; <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>).

Under the terms of this licence, you may copy, redistribute and adapt the work for non-commercial purposes, provided the work is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that WHO endorses any specific organization, products or services. The use of the WHO logo is not permitted. If you adapt the work, then you must license your work under the same or equivalent Creative Commons licence. If you create a translation of this work, you should add the following disclaimer along with the suggested citation: “This translation was not created by the World Health Organization (WHO). WHO is not responsible for the content or accuracy of this translation. The original English edition shall be the binding and authentic edition”.

Any mediation relating to disputes arising under the licence shall be conducted in accordance with the mediation rules of the World Intellectual Property Organization (<http://www.wipo.int/amc/en/mediation/rules/>).

Suggested citation. Yoong SL, Tursan d’Espaignet E, Wiggers J, St Claire S, Mellin-Olsen J, Grady A et al. Tobacco and postsurgical outcomes: WHO tobacco knowledge summaries. Geneva: World Health Organization; 2020. Licence: [CC BY-NC-SA 3.0 IGO](#).

Cataloguing-in-Publication (CIP) data. CIP data are available at <http://apps.who.int/iris>.

Sales, rights and licensing. To purchase WHO publications, see <http://apps.who.int/bookorders>. To submit requests for commercial use and queries on rights and licensing, see <http://www.who.int/about/licensing>.

Third-party materials. If you wish to reuse material from this work that is attributed to a third party, such as tables, figures or images, it is your responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned component in the work rests solely with the user.

General disclaimers. The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of WHO concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers’ products does not imply that they are endorsed or recommended by WHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by WHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall WHO be liable for damages arising from its use.

The named authors alone are responsible for the views expressed in this publication.

Acknowledgements

External reviewers: Davy Cheng,¹ Jean Wong² and Susan Ming³

¹ London Health Sciences Centre, Canada; ² University Health Network, Canada; ³ University of California San Francisco, USA

Internal WHO reviewers: Rachel Davis and Walter Johnson (WHO, Geneva).

Editorial assistance: Teresa Lander

TOBACCO & POSTSURGICAL OUTCOMES

References

1. World Health Organization (2018). WHO global report on trends in prevalence of tobacco smoking 2000–2025, second edition. Geneva: World Health Organization (<http://www.who.int/tobacco/publications/surveillance/trends-tobacco-smoking-second-edition>).
2. U.S. Department of Health and Human Services. The Health Consequences of Smoking – 50 years of Progress. A report of the Surgeon general. Atlanta, GA: U.S. Department of Health and human services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and health promotion, Office of Smoking and Health, 2014.
3. GBD 2015 Tobacco Collaborators. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the Global Burden of Disease Study 2015. *Lancet*. 2017;389(10082):1885–1906 (<https://www.ncbi.nlm.nih.gov/pubmed/28390697>, accessed 21 April 2018).
4. Chao TE, Sharma K, Mandigo M, Hagander L, Resch SC, Weiser TG, et al. Cost-effectiveness of surgery and its policy implications for global health: a systematic review and analysis. *Lancet Glob Health*. 2014;2(6):e334–e45.
5. Weiser TG, Haynes AB, Molina G, Lipsitz SR, Esquivel MM, Uribe-Leitz T, et al. Size and distribution of the global volume of surgery in 2012. *Bulletin World Health Organ*. 2016;94(3):201.
6. Weiser TG, Regenbogen SE, Thompson KD, Haynes AB, Lipsitz SR, Berry WR, et al. An estimation of the global volume of surgery: a modelling strategy based on available data. *Lancet*. 2008;372(9633):139–44.
7. Gawande AA, Thomas EJ, Zinner MJ, Brennan TA. The incidence and nature of surgical adverse events in Colorado and Utah in 1992. *Surgery*. 1999;126(1):66–75.
8. Kable AK, Gibberd RW, Spigelman AD. Adverse events in surgical patients in Australia. *Int J Qual Health*. 2002;14(4):269–76.
9. Bainbridge D, Martin J, Arango M, Cheng D. Perioperative and anaesthetic-related mortality in developed and developing countries: a systematic review and meta-analysis. *Lancet*. 2012;380(9847):1075–81.
10. Yip MK, Ng KJ. Risk-adjusted surgical audit with the POSSUM scoring system in a developing country. *Physiological and Operative Severity Score for the enUmeration of Mortality and morbidity*. The British journal of surgery. 2002;89(1):110–3.
11. McConkey SJ. Case series of acute abdominal surgery in rural Sierra Leone. *World journal of surgery*. 2002;26(4):509–13.
12. Vonlanthen R, Slankamenac K, Breitenstein S, Puhon MA, Muller MK, Hahnloser D, et al. The impact of complications on costs of major surgical procedures: a cost analysis of 1200 patients. *Annals of surgery*. 2011;254(6):907–13.
13. Moller A, Tonnesen H. Risk reduction: perioperative smoking intervention. *Best practice & research Clinical anaesthesiology*. 2006;20(2):237–48.
14. Sorensen LT. Wound healing and infection in surgery. The clinical impact of smoking and smoking cessation: a systematic review and meta-analysis. *Arch Surg*. 2012;147(4):373–83.
15. Xing D, Ma J-X, Ma X-L, Song D-H, Wang J, Chen Y, et al. A methodological, systematic review of evidence-based independent risk factors for surgical site infections after spinal surgery. *Eur Spine J*. 2013;22(3):605–15.
16. van Adrichem LN, Hovius SE, van Strik R, van der Meulen JC. Acute effects of cigarette smoking on microcirculation of the thumb. *British journal of plastic surgery*. 1992;45(1):9–11.
17. Sarin CL, Austin JC, Nickel WO. Effects of smoking on digital blood-flow velocity. *Jama*. 1974;229(10):1327–8.
18. Gourgiotis S, Aloizos S, Aravosita P, Mystakelli C, Isaia E-C, Gakis C, et al. The effects of tobacco smoking on the incidence and risk of intraoperative and postoperative complications in adults. *Surg*. 2011;9(4):225–32.
19. Tonnesen H, Nielsen PR, Lauritzen JB, Moller AM. Smoking and alcohol intervention before surgery: evidence for best practice. *Br J Anaesth*. 2009;102(3):297–306.
20. Moores LK. Smoking and postoperative pulmonary complications: An Evidence-Based Review of the Recent Literature. *Clin Chest Med*. 2000;21(1):139–46.
21. Vieira Cavichio B, Alcalá Pompeo D, de Oliveira Oller GASA, Aparecida Rossi L. Duration of smoking cessation for the prevention of surgical wound healing complications. *Rev Esc Enferm USP*. 2014;48(1):170–6.
22. Sarraf KM, Tavare A, Somashekar N, Langstaff RJ. Non-union of an undisplaced radial styloid fracture in a heavy smoker: revisiting the association of smoking and bone healing. *Hand surgery: an international journal devoted to hand and upper limb surgery and related research: journal of the Asia-Pacific Federation of Societies for Surgery of the Hand*. 2011;16(1):73–6.
23. Santolini E, West R, Giannoudis PV. Risk factors for long bone fracture non-union: a stratification approach based on the level of the existing scientific evidence. *Injury*. 2015;46 (8):S8–S19.
24. He J, Vupputuri S, Allen K, Prerost MR, Hughes J, Whelton PK. Passive Smoking and the Risk of Coronary Heart Disease — A Meta-Analysis of Epidemiologic Studies. *N Engl J Med*. 1999;340(12):920–6.
25. Boffetta P, Straif K. Use of smokeless tobacco and risk of myocardial infarction and stroke: systematic review with meta-analysis. *BMJ*. 2009;339:b3060.
26. Gronkjaer M, Eliassen M, Skov-Ettrup LS, Tolstrup JS, Christiansen AH, Mikkelsen SS, et al. Preoperative smoking status and postoperative complications: a systematic review and meta-analysis. *Annals of surgery*. 2014;259(1):52–71.
27. Wong J, Lam DP, Abrishami A, Chan MTV, Chung F. Short-term pre-operative smoking cessation and postoperative complications: a systematic review and meta-analysis. *Can J Anaesth*. 2012;59(3):268–79.
28. Mills E, Eyawo O, Lockhart I, Kelly S, Wu P, Ebbert JO. Smoking cessation reduces postoperative complications: a systematic review and meta-analysis. *Am J Med*. 2011;124(2):144–54.e8.
29. Myers K, Hajek P, Hinds C, McRobbie H. Stopping smoking shortly before surgery and postoperative complications: a systematic review and meta-analysis. *Arch Intern Med*. 2011;171(11):983–9.
30. Khullar D, Maa J. The impact of smoking on surgical outcomes. *J Am Coll Surg*. 2012;215(3):418–26.
31. Kanneganti P, Harris JD, Brophy RH, Carey JL, Lattermann C, Flanigan DC. The effect of smoking on ligament and cartilage surgery in the knee: a systematic review. *Am J Sports Med*. 2012;40(12):2872–8.
32. Singh JA. Smoking and outcomes after knee and hip arthroplasty: a systematic review. *J Rheumatol*. 2011;38(9):1824–34.
33. Reese GE, Nanidis T, Borysiewicz C, Yamamoto T, Orchard T, Tekkis PP. The effect of smoking after surgery for Crohn's disease: a meta-analysis of observational studies. *Int J Colorectal Dis*. 2008;23(12):1213–21.
34. Willigendael EM, Teijink JAW, Bartelink M-L, Peters RJG, Buller HR, Prins MH. Smoking and the patency of lower extremity bypass grafts: a meta-analysis. *J Vasc Surg*. 2005;42(1):67–74.
35. Javed F, Al-Rasheed A, Almas K, Romanos GE, Al-Hezaimi K. Effect of cigarette smoking on the clinical outcomes of periodontal surgical procedures. *Am J Med Sci*. 2012;343(1):78–84.
36. Burcharth J, Pommergaard HC, Bisgaard T, Rosenberg J. Patient-related risk factors for recurrence after inguinal hernia repair: a systematic review and meta-analysis of observational studies. *Surgical innovation*. 2015;22(3):303–17.
37. Teng S, Yi C, Krettek C, Jagodzinski M. Smoking and Risk of Prosthesis-Related Complications after Total Hip Arthroplasty: A Meta-Analysis of Cohort Studies. *PLoS ONE*. 2015;10(4):e0125294.

38. Pluvy I, Panouilleres M, Garrido I, Pauchot J, Saboye J, Chavoin JP, et al. Smoking and plastic surgery, part II. Clinical implications: a systematic review with meta-analysis. *Annales de chirurgie plastique et esthetique*. 2015;60(1):e15-49.
39. Thomsen T, Villebro N, Moller AM. Interventions for preoperative smoking cessation. *Cochrane Database Syst Rev*. 2014;3:CD002294.
40. Zaki A, Abrishami A, Wong J, Chung FF. Interventions in the preoperative clinic for long term smoking cessation: a quantitative systematic review. *Can J Anaesth*. 2008;55(1):11-21.
41. Chiswell C, Akram Y. Impact of environmental tobacco smoke exposure on anaesthetic and surgical outcomes in children: a systematic review and meta-analysis. *Arch Dis Child*. 2016(102):123-30.
42. Skolnick ET, Vomvolakis MA, Buck KA, Mannino SF, Sun LS. Exposure to Environmental Tobacco Smoke and the Risk of Adverse Respiratory Events in Children Receiving General Anesthesia. *Anesthesiology*. 1998;88(5):1144-53.
43. Seyidov TH, Elemen L, Solak M, Tugay M, Toker K. Passive smoke exposure is associated with perioperative adverse effects in children. *J Clin Anesth* [Internet]. 2011; 23(1):[47-52 pp.]. Available from: [http://www.jcafulltextonline.com/article/S0952-8180\(10\)00362-4/pdf](http://www.jcafulltextonline.com/article/S0952-8180(10)00362-4/pdf).
44. Drongowski RA, Lee D, Reynolds PI, Malviya S, Harmon CM, Geiger J, et al. Increased respiratory symptoms following surgery in children exposed to environmental tobacco smoke. *Paediatric anaesthesia*. 2003;13(4):304-10.
45. Dennis A, Curran J, Sherriff J, Kinnear W. Effects of passive and active smoking on induction of anaesthesia. *Br J Anaesth*. 1994;73(4):450-2.
46. Seyidov TH, Elemen L, Solak M, Tugay M, Toker K. Passive smoke exposure is associated with perioperative adverse effects in children. *J Clin Anesth*. 2011;23(1):47-52.
47. Simsek E, Karaman Y, Gonullu M, Tekgul Z, Cakmak M. The effect of passive exposure to tobacco smoke on perioperative respiratory complications and the duration of recovery. *Braz J Anesthesiol*. 2016;66(5):492-8.
48. Warner MA, Divertie MB, Tinker JH. Preoperative Cessation of Smoking and Pulmonary Complications in Coronary Artery Bypass Patients. *Anesthesiology*. 1984;60(4):380-3.
49. Warner MA, Offord KP, Warner ME, Lennon RL, Conover MA, Jansson-Schumacher U. Role of preoperative cessation of smoking and other factors in postoperative pulmonary complications: a blinded prospective study of coronary artery bypass patients. *Mayo Clinic proceedings*. 1989;64(6):600-16.
50. Lee SM, Landry J, Jones PM, Buhrmann O, Morley-Forster P. Long-term quit rates after a perioperative smoking cessation randomized controlled trial. *Anesth Analg*. 2015;120(3):582-7.
51. Wong J, Abrishami A, Riazi S, Siddiqui N, You-Ten E, Korman J, et al. A Perioperative Smoking Cessation Intervention With Varenicline, Counseling, and Fax Referral to a Telephone Quitline Versus a Brief Intervention: A Randomized Controlled Trial. *Anesth Analg*. 2017;125(2):571-9.
52. Rigotti NA, Clair C, Munafo MR, Stead LF. Interventions for smoking cessation in hospitalised patients. *Cochrane Database Syst Rev*. 2012;5:CD001837.
53. Stead Lindsay F, Lancaster T. Behavioural interventions as adjuncts to pharmacotherapy for smoking cessation. *Cochrane Database Syst Rev* [Internet]. 2012; 12;12:CD009670(12). Available from: <http://onlinelibrary.wiley.com/doi/10.1002/14651858.CD009670.pub2/abstract>.
54. Quraishi SA, Orkin FK, Roizen MF. The anesthesia preoperative assessment: an opportunity for smoking cessation intervention. *J Clin Anesth*. 2006;18(8):635-40.
55. Warner DO. Helping surgical patients quit smoking: time to bring it home. *Anesth Analg*. 2015;120(3):510-2.
56. Tonnesen H, Faurschou P, Ralov H, Molgaard-Nielsen D, Thomas G, Backer V. Risk reduction before surgery. The role of the primary care provider in preoperative smoking and alcohol cessation. *BMC Health Serv Res*. 2010;10:121.
57. Kotsakis GA, Javed F, Hinrichs JE, Karoussis IK, Romanos GE. Impact of cigarette smoking on clinical outcomes of periodontal flap surgical procedures: a systematic review and meta-analysis. *Journal of periodontology*. 2015;86(2):254-63.
58. Lassig AAD, Yueh B, Joseph AM. The effect of smoking on perioperative complications in head and neck oncologic surgery. *Laryngoscope*. 2012;122(8):1800-8.
59. Moller A, Villebro N. Interventions for preoperative smoking cessation. *Cochrane Database Syst Rev*. 2005;3.
60. Pluvy I, Garrido I, Pauchot J, Saboye J, Chavoin JP, Tropet Y, et al. Smoking and plastic surgery, part I. Pathophysiological aspects: Update and proposed recommendations. *Annales de chirurgie plastique et esthetique*. 2015;60(1):e15-e49.
61. Santiago-Torres J, Flanigan DC, Butler RB, Bishop JY. The effect of smoking on rotator cuff and glenoid labrum surgery: a systematic review. *Am J Sports Med*. 2015;43(3):745-51.

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_24842

