

CLIMATE AND HEALTH COUNTRY PROFILE – 2015


ETHIOPIA



World Health
Organization



United Nations
Framework Convention on
Climate Change



OVERVIEW

Ethiopia is a country with diverse climatic regions, ranging from lowland deserts and semi-arid zones to temperate zones and highland areas. Ethiopia is the second-most populous country in Sub-Saharan Africa and despite extreme poverty has experienced recent economic growth and social development [World Bank, Country Overview, 2015]. However, Ethiopia is vulnerable to many of the effects of climate change, including increases in average temperature and changes in precipitation. This threatens health, livelihoods and the progress that Ethiopia has made in recent years. Ethiopia was the first country to submit its Intended Nationally Determined Contribution to the UNFCCC and has strong plans to develop a green economy.

SUMMARY OF KEY FINDINGS

- Under a high emissions scenario, mean annual temperature is projected to rise by about 4.8°C on average from 1990 to 2100. If emissions decrease rapidly, the temperature rise is limited to about 1.3°C.
- Under a high emissions scenario, the number of days of warm spell ('heat wave') is projected to increase from about 10 days in 1990 to about 250 days on average in 2100. If emissions decrease rapidly, the days of warm spell, in 2100, are limited to about 85 on average.
- Under a high emissions scenario heat-related deaths in the elderly (65+ years) are projected to increase to over 65 deaths per 100,000 by 2080 compared to the estimated baseline of under 3 deaths per 100,000 annually between 1961 and 1990. A rapid reduction in emissions could limit heat-related deaths in the elderly to just under 12 deaths per 100,000 towards 2080.

- Under a high emissions scenario, diarrhoeal deaths attributable to climate change in children under 15 years old is projected to be about 9.6% of the over 42,000 diarrhoeal deaths projected by 2050. Although diarrhoeal deaths are projected to decline to about 15,500 towards 2070 the proportion of deaths attributable to climate change is projected to rise to approximately 14.1%.

OPPORTUNITIES FOR ACTION

Ethiopia has an approved national health adaptation strategy and has conducted a national assessment of climate change impacts, vulnerability and adaptation for health. Additionally, Ethiopia has implemented actions to build institutional and technical capacities to work on climate change and health and is currently implementing projects on health adaptation to climate change. Country reported data (see section 6) indicate there are further opportunities for action in the following areas:

1) Adaptation

- Implement activities to increase climate resilience of infrastructure including health infrastructure.
- Estimate costs to implement health resilience to climate change.

2) Mitigation

- Conduct a valuation of co-benefits of health implications of climate mitigation policies.

DEMOGRAPHIC ESTIMATES

| | |
|--|------------|
| Population [2013] ^a | 95 million |
| Population growth rate [2013] ^a | 2.5% |
| Population living in urban areas [2013] ^b | 18.6% |
| Population under five [2013] ^a | 15.1 |
| Population aged 65 or older [2013] ^a | 3.4 |

ECONOMIC AND DEVELOPMENT INDICATORS

| | |
|---|---------|
| GDP per capita [current US\$, 2013] ^c | 503 USD |
| Total expenditure on health as % of GDP [2013] ^d | 5.1% |
| Percentage share of income for lowest 20% of population [2010] ^e | 8.0% |
| HDI [2013, +/- 0.01 change from 2005 is indicated with arrow] ^e | 0.435 ▲ |

HEALTH ESTIMATES

| | |
|--|----|
| Life expectancy at birth [2013] ^f | 65 |
| Under-5 mortality per 1000 live births [2013] ^g | 65 |

a World Population Prospects: The 2015 Revision, UNDESA [2015]

b World Urbanization Prospects: The 2014 Revision, UNDESA [2014]

c World Development Indicators, World Bank [2015]

d Global Health Expenditure Database, WHO [2014]

e United Nations Development Programme, Human Development Reports [2014]

f Global Health Observatory, WHO [2014]

g Levels & Trends in Child Mortality Report 2015, UN Inter-agency Group for Child Mortality Estimation [2015]

CURRENT AND FUTURE CLIMATE HAZARDS


Due to climate change, many climate hazards and extreme weather events, such as heat waves, heavy rainfall and droughts, could become more frequent and more intense in many parts of the world.

Outlined here are country-specific projections up to the year 2100 for climate hazards under a 'business as usual' high emissions scenario compared to projections under a 'two-degree' scenario with rapidly decreasing global emissions. Most hazards caused by climate change will persist for many centuries.

COUNTRY-SPECIFIC CLIMATE HAZARD PROJECTIONS


The model projections below present climate hazards under a high emissions scenario, Representative Concentration Pathway 8.5 [RCP8.5] (in orange) and a low emissions scenario, [RCP2.6] (in green).^a The text boxes describe the projected changes averaged across about 20 models (thick line). The figures also show each model individually as well as the 90% model range (shaded) as a measure of uncertainty and, where available, the annual and smoothed observed record (in blue).^{b,c}

MEAN ANNUAL TEMPERATURE




Under a high emissions scenario, mean annual temperature is projected to rise by about 4.8°C on average from 1990 to 2100. If emissions decrease rapidly, the temperature rise is limited to about 1.3°C.

DAYS OF WARM SPELL ('HEAT WAVES')




Under a high emissions scenario, the number of days of warm spell^d is projected to increase from about 10 days in 1990 to about 250 days on average in 2100. If emissions decrease rapidly, the days of warm spell are limited to about 85 on average.

DAYS WITH EXTREME RAINFALL ('FLOOD RISK')



Under a high emissions scenario, the number of days with very heavy precipitation [20 mm or more] could double [an increase of about 6 days on average] from 1990 to 2100, increasing the risk of floods. Some models indicate increases well outside the range of historical variability, implying even greater risk. If emissions decrease rapidly, there is no increase in risk.

CONSECUTIVE DRY DAYS ('DROUGHT')



Under a high emissions scenario, the longest dry spell is indicated to decrease slightly from an average of about 95 days to just over 80 days, with continuing large year-to-year variability. If emissions decrease rapidly, the anticipated changes in the length of dry spells are somewhat less.

^a Model projections are from CMIP5 for RCP8.5 [high emissions] and RCP2.6 [low emissions]. Model anomalies are added to the historical mean and smoothed.

^b Observed historical record of mean temperature is from CRU-TSv3.22; observed historical records of extremes are from HadEX2.

^c Analysis by the Climatic Research Unit and Tyndall Centre for Climate Change Research, University of East Anglia, 2015.


^d A 'warm spell' day is a day when maximum temperature, together with that of at least the 6 consecutive previous days, exceeds the 90th percentile threshold for that time of the year.

CURRENT AND FUTURE HEALTH RISKS DUE TO CLIMATE CHANGE

Human health is profoundly affected by weather and climate. Climate change threatens to exacerbate today's health problems – deaths from extreme weather events, cardiovascular and respiratory diseases, infectious diseases and malnutrition – whilst undermining water and food supplies, infrastructure, health systems and social protection systems.

EXPOSURE TO INLAND FLOOD RISK

Population affected by inland flood risk in Ethiopia (in thousands)




Under a high emission scenario, it is projected that by 2030, 248,200 additional people may be at risk of river floods annually due to climate change and 178,100 due to socio-economic change above the estimated 154,400 annual affected population in 2010.

Source: World Resources Institute, Global Aqueduct Flood Analyzer; Assumes continued current socio-economic trends [SSP2] and a 10-year flood plan.

INFECTIOUS AND VECTOR-BORNE DISEASES


Population at risk of malaria in Ethiopia (in millions)



By 2070, almost 130 million people are projected to be at risk of malaria assuming a high emissions scenario. Population growth can also cause increases in the population at risk in areas where malaria presence is static in the future.

Source: Rocklöv, J., Quam, M. et al. 2015^b

Estimated number of deaths due to diarrhoeal disease in children under 15 years in Ethiopia, (base case scenario for economic growth)



^a Atlas of Health and Climate, World Health Organization and World Meteorological Organization, 2012.

^b Country-level analysis, completed in 2015, was based on health models outlined in the Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014.

^c WHO. Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014.



KEY IMPLICATIONS FOR HEALTH

In addition to deaths from drowning, flooding causes extensive indirect health effects, including impacts on food production, water provision, ecosystem disruption, infectious disease outbreak and vector distribution. Longer term effects of flooding may include post-traumatic stress and population displacement.

Adaptation interventions such as building climate resilient infrastructure, ecosystem-based adaptation, and efforts to strengthen community resilience can substantially reduce these figures.



KEY IMPLICATIONS FOR HEALTH

Some of the world's most virulent infections are also highly sensitive to climate: temperature, precipitation and humidity have a strong influence on the life-cycles of the vectors and the infectious agents they carry and influence the transmission of water and food-borne diseases.^a


Under a high emissions scenario, the mean relative vectorial capacity for dengue fever transmission is projected to increase to 0.57 by 2070 from 0.44 during the baseline period, remaining at a relatively high endemic transmission level towards 2070. [Source: Rocklöv, J., Quam, M. et al., 2015]^b

In 2008, there was an estimated 85,300 diarrhoeal deaths in children under 15. Under a high emissions scenario, diarrhoeal deaths attributable to climate change in children under 15 years old is projected to be about 9.6% of the over 42,000 diarrhoeal deaths projected by 2030. Although diarrhoeal deaths are projected to decline to about 15,500 towards 2050 the proportion of deaths attributable to climate change will rise to approximately 14.1%. [Source: Lloyd, S., 2015]^b

Socioeconomic development and health interventions are driving down burdens of several infectious diseases, and these projections assume that this will continue. However, climate conditions are projected to become significantly more favourable for transmission, slowing progress in reducing burdens, and increasing the populations at risk if control measures are not maintained or strengthened.^c

HEAT-RELATED MORTALITY

Heat-related mortality in population 65 years or over, Ethiopia (deaths / 100,000 population 65+ years)



Under a high emissions scenario heat-related deaths in the elderly (65+ years) are projected to increase to over 65 deaths per 100,000 by 2080 compared to the estimated baseline of under 3 deaths per 100,000 annually between 1961 and 1990. A rapid reduction in emissions could limit heat-related deaths in the elderly to just under 12 deaths per 100,000 towards 2080.

Source: Honda et al., 2015.^a




KEY IMPLICATIONS FOR HEALTH

Climate change is expected to increase mean annual temperature and the intensity and frequency of heat waves resulting in a greater number of people at risk of heat-related medical conditions.

The elderly, children, the chronically ill, the socially isolated and at-risk occupational groups are particularly vulnerable to heat-related conditions.

HEAT STRESS AND LABOUR PRODUCTIVITY

Annual daily work hours lost in relation to change in global mean temperature, Ethiopia (%)



Labour productivity is projected to decline significantly under a high emissions scenario. If global mean temperature rises 4 degrees, over 2% of annual daily work hours is projected to be lost by workers carrying out heavy labour (e.g. agricultural, construction and some industrial workers).

UNDERNUTRITION

Climate change, through higher temperatures, land and water scarcity, flooding, drought and displacement, negatively impacts agricultural production and causes breakdown in food systems. These disproportionately affect those most vulnerable to hunger and can lead to food insecurity. Vulnerable groups risk further deterioration into food and nutrition crises if exposed to extreme weather events.^b

Without considerable efforts made to improve climate resilience, it has been estimated that the risk of hunger and malnutrition globally could increase by up to 20 percent by 2050.^b

In Ethiopia, the prevalence of child malnutrition in children under age 5 is 25.2% [2014].^c

^a Country-level analysis, completed in 2015, was based on health models outlined in the Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014.

^b World Food Project 2015 <https://www.wfp.org/content/two-minutes-climate-change-and-hunger>

^c World Health Organization, Global Database on Child Growth and Malnutrition [2015 edition]. Child malnutrition estimates are for % underweight, defined as: Percentage of children aged 0-59 months who are below minus two standard deviations from median weight-for-age of the World Health Organization (WHO) Child Growth Standards.

CURRENT EXPOSURES AND HEALTH RISKS DUE TO AIR POLLUTION

Many of the drivers of climate change, such as inefficient and polluting forms of energy and transport systems, also contribute to air pollution. Air pollution is now one of the largest global health risks, causing approximately seven million deaths every year. There is an important opportunity to promote policies that both protect the climate at a global level, and also have large and immediate health benefits at a local level.

OUTDOOR AIR POLLUTION & SHORT LIVED CLIMATE POLLUTANTS



KEY IMPLICATIONS FOR HEALTH

Short-lived climate pollutants (SLCPs) such as black carbon, methane and tropospheric ozone are released through inefficient use and burning of biomass and fossil fuels for transport, housing, power production, industry, waste disposal (municipal and agricultural) and forest fires. SLCPs are responsible for a substantial fraction of global warming as well as air-pollution related deaths and diseases.

Since short lived climate pollutants persist in the atmosphere for weeks or months while CO₂ emissions

persist for years, significant reductions of SLCP emissions could reap immediate health benefits and health cost savings, and generate very rapid climate benefits – helping to reduce near-term climate change by as much as 0.5°C before 2050.^a

In Ethiopia, it is projected that a reduction in SLCPs* could prevent 24,800 premature deaths per year from outdoor air pollution (PM_{2.5} and ozone), from 2030 onwards [Source: Shindell, D., Science, 2012].

* Through implementation of 14 reduction measures: 7 targeting methane emissions and the rest, emissions from incomplete combustion. See source for further detail.


HOUSEHOLD AIR POLLUTION

ETHIOPIA

Percentage of population primarily using solid fuels for cooking (%), 2013



Source: Global Health Observatory, data repository, World Health Organization, 2013.



Source: Global Health Observatory, data repository, World Health Organization, 2012.



KEY IMPLICATIONS FOR HEALTH

Air pollution in and around the home is largely a result of the burning of solid fuels (biomass or coal) for cooking.

Women and children are at a greater risk for disease from household air pollution. Consequently, household air pollution is responsible for a larger proportion of the total number of deaths from ischaemic heart disease, stroke, lung cancer and COPD in women compared to men.^b

In Ethiopia, 63% percent of an estimated 36,800 child deaths due to acute lower respiratory infections is attributable to household air pollution [WHO, 2012].

a United Nations Environment Programme. Reducing Climate-related Air Pollution and Improving Health: Countries can act now and reap immediate benefits. <http://www.unep.org/ccac/Media/PressReleases/ReducingClimate-relatedAirPollution/tabid/131802/language/en-US/Default.aspx>

b Annu. Rev. Public. Health. 2014.35:185-206. http://www.who.int/phe/health_topics/outdoorair/databases/HAP_BoD_results_March2014.pdf?ua=1

CO-BENEFITS TO HEALTH FROM CLIMATE CHANGE MITIGATION: A GLOBAL PERSPECTIVE

Health co-benefits are local, national and international measures with the potential to simultaneously yield large, immediate public health benefits and reduce the upward trajectory of greenhouse gas emissions. Lower carbon strategies can also be cost-effective investments for individuals and societies.

Presented here are examples, from a global perspective, of opportunities for health co-benefits that could be realised by action in important greenhouse gas emitting sectors.^a

Transport

Transport injuries lead to 1.2 million deaths every year, and land use and transport planning contribute to the 2–3 million deaths from physical inactivity. The transport sector is also responsible for some 14% (7.0 GtCO₂e) of global carbon emissions. The IPCC has noted significant opportunities to reduce energy demand in the sector, potentially resulting in a 15%–40% reduction in CO₂ emissions, and bringing substantial opportunities for health: A modal shift towards walking and cycling could see reductions in illnesses related to physical inactivity and reduced outdoor air pollution and noise exposure; increased use of public transport is likely to result in reduced GHG emissions; compact urban planning fosters walkable residential neighborhoods, improves accessibility to jobs, schools and services and can encourage physical activity and improve health equity by making urban services more accessible to the elderly and poor.



Electricity Generation

Reliable electricity generation is essential for economic growth, with 1.4 billion people living without access to electricity. However, current patterns of electricity generation in many parts of the world, particularly the reliance on coal combustion in highly polluting power plants contributes heavily to poor local air quality, causing cancer, cardiovascular and respiratory disease. Outdoor air pollution is responsible for 3.7 million premature deaths annually, 88% of these deaths occur in low and middle income countries. The health benefits of transitioning from fuels such as coal to lower carbon sources, including ultimately to renewable energy, are clear: Reduced rates of cardiovascular and respiratory disease such as stroke, lung cancer, coronary artery disease, and COPD; cost-savings for health systems; improved economic productivity from a healthier and more productive workforce.



Household Heating, Cooking and Lighting

Household air pollution causes over 4.3 million premature deaths annually, predominantly due to stroke, ischaemic heart disease, chronic respiratory disease, and childhood pneumonia. A range of interventions can both improve public health and reduce household emissions: a transition from the inefficient use of solid fuels like wood and charcoal, towards cleaner energy sources like liquefied petroleum gas (LPG), biogas, and electricity could save lives by reducing indoor levels of



Healthcare Systems

Health care activities are an important source of greenhouse gas emissions. In the US and in EU countries, for example, health care activities account for between 3–8% of greenhouse gas (CO₂-eq) emissions. Major sources include procurement and inefficient energy consumption. Modern, on-site, low-carbon energy solutions (e.g. solar, wind, or hybrid solutions) and the development of combined heat and power generation capacity in larger facilities offer significant potential to lower the health sector's carbon footprint.



预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_26727

