## CLIMATE AND HEALTH COUNTRY PROFILE - 2015 **BHUTAN**





**United Nations** Framework Convention on Climate Change



### **OVERVIEW**

The Kingdom of Bhutan in the eastern Himalayas is a mountainous, landlocked country with 70% forest cover [Bhutan INDC, 2015]. Climate varies with elevation and season with subtropical plains in the south and year-round snow in the Himalayan heights of the north.

Bhutan is recognized for its measure of Gross National Happiness over GDP, its commitment to sustainable development and being a net carbon sink - taking in more carbon dioxide than it produces. Bhutan has pledged to remain carbon neutral and has a constitutional mandate to achieve this by maintaining forest cover at over 60% [Bhutan INDC, 2015]. Economic growth has been largely driven by the hydropower sector [World Bank Country Overview, 2015].

Despite positive national action on climate change, Bhutan is vulnerable to climate change and extreme events, most significantly glacial lake outburst floods but also flash floods and forest fires. Additional health concerns include an increase in the geographic range and incidence of vector-borne diseases such as dengue fever, and the incidence of water-borne diseases.

### SUMMARY OF KEY FINDINGS

- In Bhutan, under a high emissions scenario, mean annual temperature is projected to rise by about 5.6°C on average from 1990 to 2100. If global emissions decrease rapidly, the temperature rise is limited to about 1.6°C (page 2).
- In Bhutan, under a high emissions scenario heat-related deaths in the elderly (65+ years) are projected to increase

to about 49 deaths per 100,000 by 2080 compared to the estimated baseline of zero deaths per 100,000 annually between 1961 and 1990. A rapid reduction in global emissions could limit heat-related deaths in the elderly to about 8 deaths per 100,000 in 2080 (page 4).

 Labour productivity, in Bhutan, is projected to decline under a high emissions scenario. If global mean temperature rises 4 degrees, about 12% of annual daily work hours is projected to be lost by workers carrying out heavy labour (page 4).

### **OPPORTUNITIES FOR ACTION**

Bhutan has an approved national health adaptation strategy and is currently implementing projects on health adaptation to climate change. Bhutan has conducted a national assessment of climate change impacts, vulnerabilities and adaptation for health. Country reported data (see section 6) indicate that there remains opportunities for action in the following areas:

### 1) Adaptation

- Implement actions to build technical and institutional capacities to work on climate change and health.
- Implement activities to increase climate resilience of health
- · Estimate the costs of implementing health resilience to climate change.

#### 2) Mitigation

 Conduct a valuation of co-benefits to health of climate change mitigation policies.

DEMOGRAPHIC ESTIMATES	
Population (2013) <sup>a</sup>	0.76 million
Population growth rate (2013) <sup>a</sup>	1.4 %
Population living in urban areas (2013) <sup>b</sup>	37.1 %
Population under five (2013) <sup>a</sup>	9.1 %
Population aged 65 or older (2013)ª	4.8 %
ECONOMIC AND DEVELOPMENT INDICATORS	
GDP per capita (current US\$, 2013) <sup>c</sup>	2,383 USD
Total expenditure on health as % of GDP (2013) <sup>d</sup>	3.6 %
Percentage share of income for lowest 20% of population (2012) <sup>c</sup>	6.8
HDI (2013) <sup>e</sup>	0.584
HEALTH ESTIMATES	
Life expectancy at birth (2013) <sup>f</sup>	68 years
Under-5 mortality per 1000 live births (2013) <sup>9</sup>	36

World Urbanization Prospects: The 2014 Revision, UNDESA [2014] World Development Indicators, World Bank [2015] Global Health Expenditure Database, WHO [2014]

- United Nations Development Programme, Human Development Reports [2014] Global Health Observatory, WHO [2014] Levels & Trends in Child Mortality Report 2015, UN Inter-agency Group for Child Mortality Estimation [2015]

## 1 CURRENT AND FUTURE CLIMATE HAZARDS


Due to climate change, many climate hazards and extreme weather events, such as heat waves, heavy rainfall and droughts, could become more frequent and more intense in many parts of the world.

Outlined here are country–specific projections up to the year 2100 for climate hazards under a 'business as usual' high emissions scenario compared to projections under a 'two-degree' scenario with rapidly decreasing global emissions. Most hazards caused by climate change will persist for many centuries.

### **COUNTRY-SPECIFIC CLIMATE HAZARD PROJECTIONS**


The model projections below present climate hazards under a high emissions scenario, Representative Concentration Pathway 8.5 [RCP8.5] (in orange) and a low emissions scenario, [RCP2.6] (in green).<sup>a</sup> The text boxes describe the projected changes averaged across about 20 models (thick line). The figures also show each model individually as well as the 90% model range (shaded) as a measure of uncertainty and, where available, the annual and smoothed observed record (in blue).<sup>b,c</sup>

### **MEAN ANNUAL TEMPERATURE**




Under a high emissions scenario, mean annual temperature is projected to rise by about  $5.6^{\circ}$ C on average from 1990 to 2100. If emissions decrease rapidly, the temperature rise is limited to about  $1.6^{\circ}$ C.

### DAYS WITH EXTREME RAINFALL ('FLOOD RISK')




Under a high emissions scenario, the number of days with very heavy precipitation (20 mm or more) could increase by about 5 days on average from 1990 to 2100, increasing the risk of floods. Some models indicate increases well outside the range of historical variability, implying even greater risk. If emissions decrease rapidly, the increase in risk is much reduced.

### DAYS OF WARM SPELL ('HEAT WAVES')



Under a high emissions scenario, the number of days of warm spell<sup>d</sup> is projected to increase from less than 20 days in 1990 to about 235 days on average in 2100. If emissions decrease rapidly, the days of warm spell are limited to about 65 on average.

### **CONSECUTIVE DRY DAYS ('DROUGHT')**



Under a high emissions scenario, the longest dry spell is indicated to increase by almost 20 days from an average of about 80-90 days, with continuing large year-to-year variability. If emissions decrease rapidly, the anticipated changes in the length of dry spells are somewhat less.

- a Model projections are from CMIP5 for RCP8.5 (high emissions) and RCP2.6 (low emissions). Model anomalies are added to the historical mean and smoothed.
- b Observed historical record of mean temperature is from CRU-TSv.3.22; observed historical records of extremes are from HadEX2.
- c Analysis by the Climatic Research Unit and Tyndall Centre for Climate Change Research, University of East Anglia, 2015.
- d A 'warm spell' day is a day when maximum temperature, together with that of at least the 6 consecutive previous days, exceeds the 90th percentile threshold for that time of the year.

2


2

## CURRENT AND FUTURE HEALTH RISKS DUE TO CLIMATE CHANGE

Human health is profoundly affected by weather and climate. Climate change threatens to exacerbate today's health problems – deaths from extreme weather events, cardiovascular and respiratory diseases, infectious diseases and malnutrition – whilst undermining water and food supplies, infrastructure, health systems and social protection systems.

### **EXPOSURE TO INLAND RIVER FLOODING**

Population affected by inland river flood risk in Bhutan



It is projected, that by 2030, under a high emissions scenario, an additional 7,600 people may be at risk of river floods annually as a result of climate change and 2,400 due to socio-economic change above the estimated 3,100 annually affected population in 2010.<sup>a</sup>

### INFECTIOUS AND VECTOR-BORNE DISEASES



## **KEY IMPLICATIONS FOR HEALTH**

Rising temperatures in Bhutan and the subsequent melting of glaciers is also leading to a significant threat of flash flooding due to glacial lake outburts.<sup>b</sup>

In addition to deaths from drowning, flooding causes extensive indirect health effects, including impacts on food production, water provision, ecosystem disruption, infectious disease outbreak and vector distribution. Longer term effects of flooding may include post-traumatic stress and population displacement.

## KEY IMPLICATIONS FOR HEALTH

Some of the worlds most virulent infections are also highly sensitive to climate: temperature , precipitation and humidity have a strong influence on the life-cycles of the vectors and the infectious agents they carry and influence the transmission of water and food-borne diseases.<sup>c</sup>


Socioeconomic development and health interventions are driving down burdens of several infectious diseases, and these projections assume that this will continue. However, climate conditions are projected to become significantly more favourable for transmission, slowing progress in reducing burdens, and increasing the populations at risk if control measures are not maintained or strengthened.<sup>d</sup>

- c Atlas of Health and Climate, World Health Organization and World Meteorological Organization, 2012.
- d Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s. Geneva: World Health Organization, 2014.

a World Resources Institute, http://www.wri.org. Aqueduct Global Flood Analyzer. Assumes continued current socio-economic trends (SSP2) and a 25-year flood protection.
b UNDP in Bhutan, Climate Change Adaptation and Disaster Risk reduction, 2015. http://www.undp.org/content/bhutan/en/home/ourwork/ccaadaptationanddrr/successstories/ glofprevention.html

### **HEAT-RELATED MORTALITY**

Heat-related mortality in population 65 years or over, Bhutan (deaths / 100,000 population 65+ yrs)



Under a high emissions scenario heat-related deaths in the elderly (65+ years) are projected to increase to about 49 deaths per 100,000 by 2080 compared to the estimated baseline of zero deaths per 100,000 annually between 1961 and 1990. A rapid reduction in global emissions could limit heat-related deaths in the elderly to about 8 deaths per 100,000 in 2080.

Source: Honda et al., 2015.ª


## **KEY IMPLICATIONS FOR HEALTH**

Climate change is expected to increase mean annual temperature and the intensity and frequency of heat waves resulting in a greater number of people at risk of heat-related medical conditions.

The elderly, children, the chronically ill, the socially isolated and at-risk occupational groups are particularly vulnerable to heat-related conditions.

### HEAT STRESS AND LABOUR PRODUCTIVITY

Annual daily work hours lost in relation to change in global mean temperature, Bhutan (%)



Labour productivity is projected to decline under a high emissions scenario. If global mean temperature rises 4 degrees, about 12% of annual daily work hours is projected to be lost by workers carrying out heavy labour (e.g. agricultural, construction and some industrial workers).

Source: Kiellstrom et al., 2015 http://www.climatechip.org/

### UNDERNUTRITION

Climate change, through higher temperatures, land and water scarcity, flooding, drought and displacement, negatively impacts agricultural production and causes breakdown in food systems. These disproportionally affect those most vulnerable people at risk to hunger and can lead to food insecurity. Vulnerable groups risk further deterioration into food and nutrition crises if exposed to extreme climate events.<sup>b</sup>

Without considerable efforts made to improve climate resilience, it has been estimated that the global risk of hunger and malnutrition could increase by up to 20 percent by 2050.<sup>b</sup>

In Bhutan, the prevalence of stunting in children under age 5 was 33.6% in 2010, the prevalence of underweight children and wasting in children under 5 was 12.8% and 5.9%, respectively, in 2010.°

World Food Project 2015 https://www.wfp.org/content/two-minutes-climate-change-and-hunger World Health Organization, Global Database on Child Growth and Malnutrition [2015 edition]. Please see source for definitions of child malnutrition measures.

4


Country-level analysis, completed in 2015, was based on health models outlined in the Quantitative risk assessment of the effects of climate change on selected causes of а death, 2030s and 2050s. Geneva: World Health Organization, 2014. The mean of impact estimates for three global climate models are presented. Models assume continued socioeconomic trends (SSP2 or comparable).

## CURRENT EXPOSURES AND HEALTH RISKS DUE TO AIR POLLUTION

Many of the drivers of climate change, such as inefficient and polluting forms of energy and transport systems, also contribute to air pollution. Air pollution is now one of the largest global health risks, causing approximately seven million deaths every year. There is an important opportunity to promote policies that both protect the climate at a global level, and also have large and immediate health benefits at a local level.

### **OUTDOOR AIR POLLUTION EXPOSURE**

Outdoor air pollution in Thimpu, Bhutan annual mean PM<sub>2.5</sub> (µg/m<sup>3</sup>) 2011\*





## KEY IMPLICATIONS FOR HEALTH

Outdoor air pollution can have direct and sometimes severe consequences for health.

Fine particles which penetrate deep into the respiratory tract subsequently increase mortality from respiratory infections, lung cancer and cardiovascular disease.

In Thimpu, air pollution data in 2011\* indicated an annual mean PM<sub>2.5</sub> level that was at the WHO guideline value of 10 µg/m<sup>3</sup>.

Source: Ambient Air Pollution Database, WHO, May 2014.

\*A standard conversion has been used, see source for further details.

### HOUSEHOLD AIR POLLUTION

### **BHUTAN**


3

Percentage of population primarily using solid fuels for cooking (%), 2013



Source: Global Health Observatory, data repository, World Health Organization, 2013.

Percent of total deaths from ischaemic heart disease, stroke, lung cancer, chronic obstructive pulmonary disease (18 years +) and acute lower respiratory infections (under 5 years) attributable to household air pollution, 2012.





**URBAN** 

AREAS

Air pollution in and around the home is largely a result of the burning of solid fuels (biomass or coal) for cooking.

ΝΔΤΙΟΝΔΙ

TOTAL

Women and children are at a greater risk for disease from household air pollution. Consequently, household air pollution is responsible for a larger proportion of the of total number of deaths from ischaemic heart disease, stroke, lung cancer and COPD in women compared to men.<sup>a</sup>

In Bhutan, 39% percent of an estimated 110 child deaths due to acute lower respiratory infections is attributable to household air pollution (WHO, 2012).

Source: Global Health Observatory, data repository, World Health Organization, 2012.

a Annu. Rev. Public. Health. 2014.35:185-206. http://www.who.int/phe/health\_topics/outdoorair/databases/HAP\_BoD\_results\_March2014.pdf?ua=1

## 4

## CO-BENEFITS TO HEALTH FROM CLIMATE CHANGE MITIGATION: A GLOBAL PERSPECTIVE

Health co-benefits are local, national and international measures with the potential to simultaneously yield large, immediate public health benefits and reduce the upward trajectory of greenhouse gas emissions. Lower carbon strategies can also be cost-effective investments for individuals and societies.

Presented here are examples, from a global perspective, of opportunities for health co-benefits that could be realised by action in important greenhouse gas emitting sectors.<sup>a</sup>

### Transport

Transport injuries lead to 1.2 million deaths every year, and land use and transport planning contribute to the 2-3 million deaths from physical inactivity. The transport sector is also responsible for some 14% (7.0 GtCO₂e) of global carbon emissions. The IPCC has noted significant opportunities to reduce energy demand in the sector, potentially resulting in a 15%–40% reduction in CO<sub>2</sub> emissions, and bringing substantial opportunities for health: A modal shift towards walking and cycling could see reductions in illnesses related to physical inactivity and reduced outdoor air pollution and noise exposure; increased use of public transport is likely to result in reduced GHG emissions; compact urban planning fosters walkable residential neighborhoods, improves accessibility to jobs, schools and services and can encourage physical activity and improve health equity by making urban services more accessible to the elderly and poor.

### **Electricity Generation**

Reliable electricity generation is essential for economic growth, with 1.4 billion people living without access to electricity. However, current patterns of electricity generation in many parts of the world, particularly the reliance on coal combustion in highly polluting power plants contributes heavily to poor local air quality, causing cancer, cardiovascular and respiratory disease. Outdoor air pollution is responsible for 3.7 million premature deaths annually, 88% of these deaths occur in low and middle income countries. The health benefits of transitioning from fuels such as coal to lower carbon sources, including ultimately to renewable energy, are clear: Reduced rates of cardiovascular and respiratory disease such as stroke, lung cancer, coronary artery

disease, and COPD; cost-savings for health systems; improved economic productivity from a healthier and more productive workforce.

### Household Heating, Cooking and Lighting

Household air pollution causes over 4.3 million premature deaths annually, predominantly due to stroke, ischaemic heart disease, chronic respiratory disease, and childhood pneumonia. A range of interventions can both improve public health and reduce household emissions: a transition from the inefficient use of solid fuels like wood and charcoal, towards cleaner energy sources like liquefied petroleum gas (LPG), biogas, and electricity could save lives by reducing indoor levels of

### **Healthcare Systems**

Health care activities are an important source of greenhouse gas emissions. In the US and in EU countries, for example, health care activities account for between 3–8% of greenhouse gas (CO<sub>2</sub>-eq) emissions. Major sources include procurement

and inefficient energy consumption. Modern, on-site, low-carbon energy solutions (e.g. solar, wind, or hybrid solutions) and the development of combined heat and power generation capacity in larger facilities offer significant potential to lower the health sector's carbon

# 预览已结束, 完整报告链接和二维码如下:

https://www.yunbaogao.cn/report/index/report?reportId=5 26839

