



Basic Methods for
**Assessment of
Renal Fluoride
Excretion**

in Community
Prevention
Programmes for
Oral Health



World Health
Organization

**Basic methods for
assessment of renal fluoride
excretion in community
prevention programmes for
oral health**

2014



**World Health
Organization**

WHO Library Cataloguing-in-Publication Data

Basic methods for assessment of renal fluoride excretion in community prevention programmes for oral health.

1. Fluoride – adverse effects. 2. Fluoride – metabolism. 3. Fluoridation. 4. Fluorosis, Dental – prevention and control. 5. Community Dentistry. 6. Oral Health. 7. Program Evaluation – methods. I. World Health Organization.

ISBN 978 92 4 154870 0

(NLM classification: WU 270)

© **World Health Organization 2014**

All rights reserved. Publications of the World Health Organization are available on the WHO web site (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications – whether for sale or for non-commercial distribution – should be addressed to WHO Press through the WHO web site (www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

PRINTED IN FRANCE

Contents

Acronyms and abbreviations	1
Preface	2
Acknowledgements	5
1 Introduction	6
1.1 Background	6
1.2 Sources of fluoride intake in humans	7
1.3 Fluoride metabolism and excretion	8
1.4 Biomarkers of fluoride exposure	8
1.5 Urinary fluoride assessments in population oral health	9
1.6 Areas of uncertainty	10
2 General design of study	12
2.1 Background	12
2.2 Identification of participants	12
2.3 Sampling	12
2.4 Methods of monitoring	15
2.4.1 24-hour urine collection	15
2.4.2 Timed collections of urine obtained from defined periods of a day	17
3 Methods for collecting urine, and for handling and evaluating results	20
3.1 Recording of information	20
3.1.1 General information	20
3.1.2 Personal information	23
3.2 Essential preparatory practices	29
3.2.1 Pre-collection planning approaches	29
4 Twenty-four hour urine collection	33
4.1 Procedure	33
4.2 Fluoride and creatinine concentrations	34
4.3 Information to be recorded	34

4.4 Determination of fluoride in urine	35
4.5 General rules for tabulation and processing of data	35
5 Design of the final report for 24-hour urine collection	37
5.1 Introduction	37
5.2 Materials and methods	37
5.3 Results	37
5.4 Discussion and conclusion	39
5.5 Summary	39
5.6 Tables for the 24-hour collection method	40
6 Collections of 8–18 hours (within the 24-hour cycle)	44
6.1 Collection of nocturnal urine and during periods of high excretion	45
6.2 Optional calculations obtained from time-controlled urine collection – Use of a standardized format	46
6.2.1 Coded recordings of personal data and individual fluoride exposure	47
6.2.2 Standard table for surveys with 24-hour collections	48
6.2.3 Standard table for surveys with two collections totalling 14–16 hours, using MS Excel	50
6.2.4 Standard table for surveys with three collections totalling 14–16 hours using MS Excel: complete series	51
6.2.5 Standard table for surveys with collections made in four periods totalling the 24-cycle	51
6.2.6 Cleaning of data	51
6.3 Extrapolations	53
7 Design of the final report for time-controlled urinary collections	54
7.1 Tables for time-controlled urinary collections lasting 8–18 hours	55
8 Urine spot samples	65
8.1 Creatinine	65
9 Provision of technical advice	67
Annex A: Apparatus and materials	68
Annex B: Steps for verifying function of apparatus	72
Annex C: Calibration procedure	74
References	79

Tables

Table 5.1	Urinary fluoride excretion data from 24-hour urine collections	38
Table 5.2	Criteria for cleaning data from children aged 2–6 years, from 24-hour collections	40
Table 5.3	Standards for urinary fluoride excretion (mg/24-hour cycle): lower and upper margins	41
Table 5.4	Calculated daily urinary fluoride excretion associated with low, optimal and high fluoride intake for ages 1–14 years	42
Table 5.5	Calculated daily urinary fluoride excretion associated with low, optimal and high fluoride intake for broader age groups	43
Table 7.1	Urinary fluoride data for time-controlled urine collections	55
Table 7.2	Criteria for cleaning data from children aged 2–6 years, from time-controlled urine collections ^a	56
Table 7.3	Range of optimal urinary fluoride excretion ($\mu\text{g}/\text{hour}$) at different times, for time-controlled urine collections: lower and upper margins	57
Table 7.4	Standards for urinary fluoride concentration (mg/L; ppm) for all ages, for time-controlled urine collections; lower and upper margins	57
Table 7.5	Calculation of parameters using simple numerical operations	58
Table 7.6	Data and computing table using MS Excel for surveys for 24-hour collections (example with six cases)	59
Table 7.7	Dataset example and MS Excel computing table showing calculations of fluoride excretion of children providing nocturnal and noon urine collections	60
Table 7.8	Dataset example and MS Excel computing table showing calculations of fluoride excretion of children providing morning, afternoon and nocturnal collections	62
Table 7.9	Ideal format of a dataset to be transferred to an evaluation centre	64

Figures

Figure 3.1	Example of daytime label used in time-controlled collections, to be attached to the urine collecting jar (WHO Form No. 96392, modified)	21
Figure 3.2	Overnight urine collection label used in time-controlled collections (WHO Form No. 96393)	21

Basic Methods for Assessment of Renal Fluoride Excretion

Figure 3.3 Summary record form of urine collection from a group of children in time-controlled collections (WHO Form No. 96391)	22
Figure 3.4 Sample invitation	23
Figure 3.5 Questionnaire on use of fluoride supplements and toothpaste	24

Acronyms and abbreviations

bw	body weight
CV	coefficient of variation
CDTA	1,2-cyclohexylenedinitrilotetraacetic acid
DUFE	daily urinary fluoride excretion
F	fluoride
F.W.	formula weight
HPLC	high-performance liquid chromatography
ID	identification
IRB	institutional review boards
PABA	para-amino benzoic acid
ppm	parts per million
SD	standard deviation
TISAB	total ionic strength adjusting buffer
WHA	World Health Assembly
WHO	World Health Organization

预览已结束，完整报告链接和工

<https://www.yunbaogao.cn/report/index/report?re>