

WHO TobLabNet
Official Method
SOP 01

STANDARD OPERATING PROCEDURE FOR INTENSE SMOKING OF CIGARETTES

**Tobacco Free Initiative
Tobacco Laboratory Network (TobLabNet)**



**World Health
Organization**

**WHO TobLabNet
Official Method
SOP 01**

**Standard operating procedure for
intense smoking of cigarettes**

WHO Library Cataloguing-in-Publication Data

Standard operating procedure for intense smoking of cigarettes.

1.Smoking. 2.Smoke – analysis. 3.Tobacco smoke pollution – analysis.
4.Consumer product safety. 5.Toxicity tests – standards. I.World Health Organization. Tobacco Laboratory Network.

ISBN 978 92 4 150389 1

(NLM classification: WA 754)

© **World Health Organization 2012**

All rights reserved. Publications of the World Health Organization are available on the WHO web site (www.who.int) or can be purchased from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int).

Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press through the WHO web site (http://www.who.int/about/licensing/copyright_form/en/index.html).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

Printed in Switzerland

No.: SOP 01
Date: April 2012



**World Health Organization
Tobacco Laboratory Network**

**Standard operating procedure for method
Intense smoking of cigarettes**

Method:	Intense smoking of cigarettes
Analytes:	Not applicable
Matrix:	Cigarettes
Last update:	April 2012



No machine smoking regimen can represent all human smoking behaviour: machine smoking testing is useful for characterizing cigarette emissions for design and regulatory purposes, but communication of machine measurements to smokers can result in misunderstanding about differences between brands in exposure and risk. Data on smoke emissions from machine measurements may be used as inputs for product hazard assessment, but they are not intended to be nor are they valid as measures of human exposure or risks. Representing differences in machine measurements as differences in exposure or risk is a misuse of testing with WHO TobLabNet standards.

No.: SOP 01

Date: April 2012

FOREWORD

This document was prepared by members of the World Health Organization (WHO) Tobacco Laboratory Network (TobLabNet) as a standard operating procedure (SOP) for intense smoking of cigarettes.

INTRODUCTION

In order to establish comparable measurements for testing tobacco products globally, consensus methods are required for measuring specific contents and emissions of cigarettes. The Conference of the Parties to the WHO Framework Convention on Tobacco Control (FCTC) at its third session in Durban, South Africa, in November 2008, recalling its decisions FCTC/COP1(15) and FCTC/COP2(14) on the elaboration of guidelines for implementation of Articles 9 (*Regulation of the contents of tobacco products*) and 10 (*Regulation of tobacco product disclosures*) of the WHO FCTC, noting the information contained in the report of the working group to the third session of the Conference of the Parties on the progress of its work ... requested the Convention Secretariat to invite WHO's Tobacco Free Initiative to ... validate, within five years, the analytical chemical methods for testing and measuring cigarette contents and emissions (FCTC/COP/3/REC/1).

Using the criteria for prioritization set at its third meeting in Ottawa, Canada, in October 2006, the working group on Articles 9 and 10 identified the following contents for which methods for testing and measurement (analytical chemistry) should be validated as a priority:

- nicotine
- ammonia
- humectants (propane-1,2-diol, glycerol (propane-1,2,3-triol) and triethylene glycol (2,2-ethylenedioxybis(ethanol))).

Measurement of these contents will require validation of three methods: one for nicotine, one for ammonia and one for humectants.

Using the criteria for prioritization set at the meeting in Ottawa mentioned above, the working group identified the following emissions in mainstream smoke for which methods for testing and measurement (analytical chemistry) should be validated as a priority:

- 4-(methylnitrosamino)-1-(3-pyridyl)-1-butanone (NNK)
- N-nitrosornicotine (NNN)
- acetaldehyde
- acrylaldehyde (acrolein)
- benzene
- benzo[a]pyrene
- 1,3-butadiene
- carbon monoxide
- formaldehyde

Measurement of these emissions with the two smoking regimens described below will require validation of five methods: one for tobacco-specific

nitrosamines (NNK and NNN), one for benzo[a]pyrene, one for aldehydes (acetaldehyde, acrolein and formaldehyde), one for volatile organic compounds (benzene and 1,3-butadiene), and one for carbon monoxide.

The table below sets out the two smoking regimens for validation of the test methods referred to above.

Smoking regimen	Puff volume (ml)	Puff frequency	Filter ventilation holes
ISO regimen: ISO 3308; <i>Routine analytical cigarette smoking machine—definitions and standard conditions</i>	35	Once every 60 s	No modifications
Intense regimen: Same as ISO 3308, but modified as indicated	55	Once every 30 s	All ventilation holes must be blocked 100% as described in 12.2 .

This SOP was prepared to describe the procedure for intense smoking of cigarettes.

1 SCOPE

This SOP describes the overall procedures for machine smoking of cigarettes under intense conditions.

Note: Training in use of the smoking machine and other analytical equipment is important for successful operation. People not experienced in operating smoking machines or in using the analytical methods for measuring tobacco product emissions and contents should be trained.

2 REFERENCES

- 2.1 *ISO 3308: Routine analytical cigarette-smoking machine—Definitions and standard conditions.*
- 2.2 *ISO 4387: Cigarettes—Determination of total and nicotine-free dry particulate matter using a routine analytical smoking machine.*
- 2.3 *ISO 3402: Tobacco and tobacco products—Atmosphere for conditioning and testing.*

3 TERMS AND DEFINITIONS

- 3.1 TPM: Total particulate matter

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_28543

