



World Health
Organization

Global estimate of the burden of disease from second-hand smoke



Global estimate of the burden of disease from second-hand smoke

Mattias Öberg^a, Alistair Woodward^b, Maritta S. Jaakkola^c,
Armando Peruga^d, Annette Prüss-Ustün^e

^a Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden

^b School of Population Health, University of Auckland, Auckland, New Zealand

^c Respiratory Medicine Unit, Centre for Environmental and Respiratory Health research
and Institute of Clinical Medicine, University of Oulu, Oulu, Finland

^d Tobacco Free Initiative, World Health Organization, Geneva, Switzerland

^e Public Health and the Environment Department, World Health Organization, Geneva, Switzerland

WHO Library Cataloguing-in-Publication Data

Global estimate of the burden of disease from second-hand smoke / by Mattias Öberg ... [et al].

1.Smoking - epidemiology. 2.Tobacco smoke pollution - adverse effects. 3.Risk factors. 4.Cost of illness. 5.Air pollution, Indoor. I.Öberg, M. II.Jaakkola, M.S. III.Prüss-Üstün, Annette. IV.Peruga, Armando. V.Woodward, Alistair. VI.World Health Organization.

ISBN 978 92 4 156407 6

(NLM classification: HD 9130.6)

© World Health Organization 2010

All rights reserved. Publications of the World Health Organization can be obtained from WHO Press, World Health Organization, 20 Avenue Appia, 1211 Geneva 27, Switzerland (tel.: +41 22 791 3264; fax: +41 22 791 4857; e-mail: bookorders@who.int). Requests for permission to reproduce or translate WHO publications – whether for sale or for noncommercial distribution – should be addressed to WHO Press, at the above address (fax: +41 22 791 4806; e-mail: permissions@who.int).

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the World Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.

The named authors alone are responsible for the views expressed in this publication.

Printed in Switzerland

Content

ACKNOWLEDGEMENTS.....	VI
LIST OF ABBREVIATIONS.....	VII
EXECUTIVE SUMMARY	VIII
1 INTRODUCTION.....	2
1.2 DEFINING SECOND-HAND SMOKE	2
1.3 PROPERTIES OF SECOND-HAND SMOKE	2
1.4 CURRENT AND EX-SMOKERS.....	2
2 METHODS.....	3
2.1 SUMMARY OF THE METHOD USED TO ESTIMATE THE ENVIRONMENTAL BURDEN OF DISEASE AT GLOBAL LEVEL	3
2.2 EXPOSURE ASSESSMENT.....	5
2.3 HEALTH EFFECTS SELECTED FOR THE GLOBAL ESTIMATE OF BURDEN OF DISEASE	12
2.4 EQUIVALENCE OF EXPOSURE DEFINITION IN SURVEYS AND IN EPIDEMIOLOGICAL STUDIES	19
2.5 AVAILABLE HEALTH DATA FOR THE SELECTED OUTCOMES.....	26
3 RESULTS.....	31
3.1 BURDEN OF DISEASE ATTRIBUTABLE TO SECOND-HAND SMOKE	31
3.2 ESTIMATES OF UNCERTAINTY	41
3.3 SENSITIVITY ANALYSIS	43
3.4 COMPARISONS WITH OTHER STUDIES.....	44
3.5 CONCLUSIONS	46
4 REFERENCES	48
5 ANNEX.....	55
Table A1 Available national data on occurrence of second-hand smoke exposure among children and adults	55
Table A2 Estimated recent exposure to second-hand smoke for children and adult men and women.....	63
TABLES	
Table 1 Proportion of countries and children in countries covered by surveys on parental smoking, by region.....	6
Table 2 Models used for estimating children’s SHS exposure.....	7
Table 3 Proportion of children under 15 years with one or more parent who smokes (based on survey data and modelling)	9
Table 4 Proportion of adults in countries covered by surveys on exposure to SHS, by region.....	10
Table 5 The proportion of non-smoking adults exposed regularly to SHS, by WHO region, based on survey data and modelling	12
Table 6 Recommended risk estimates and summary of conclusions of recent reviews of health effects of exposure to SHS in children ^a	13
Table 7 Recommended risk estimates and summary of conclusions of recent reviews of health effects of exposure to SHS in adult non-smokers ^a	16
Table 8 Mean number of daily cigarettes consumed by smoker in each region ^a	21

Table 9	Smoking at home in the presence of children compared to parental smoking.....	22
Table 10	Hair nicotine in children and women	24
Table 11	Burden of lower respiratory infections (deaths and DALYsa) in children 0–4 years by WHO subregion, 2004	26
Table 12	Burden of acute otitis media (deaths and DALYsa) in children 0–4 years, by WHO subregion, 2004	27
Table 13	Burden of asthma (deaths and DALYsa) in children 0–14 years, by WHO subregion, 2004 ...	28
Table 14	Burden of asthma (deaths and DALYsa), by WHO subregion, in the population >15 years, 2004	29
Table 15	Burden of lung cancer (deaths and DALYsa), by WHO subregion, in the population >15 years, 2002	30
Table 16	Burden of ischaemic heart disease (deaths and DALYsa), by WHO subregion, in the population >15 years, 2004	31
Table 17	Burden of lower respiratory infections attributable to SHS exposure, in children 0–4 years, 2004.....	32
Table 18	Burden of acute otitis media attributable to SHS exposure, in children 0–4 years old, 2004.	33
Table 19	Burden of asthma attributable to SHS exposure, for children 0–14 years old, 2004	34
Table 20	Burden of asthma attributable to SHS exposure, population ≥15 years, 2004.....	35
Table 21	Burden of lung cancer attributable to SHS exposure, population ≥15 year olds, 2004 ^a	36
Table 22	Burden of ischaemic heart disease attributable to SHS exposure, opulation ≥15 years, 2004	37
Table 23	Total deaths and DALYsa from second-hand smoke by outcome, 2004	38
Table 24	Total deaths and DALYsa from second-hand smoke by subregion, 2004	38
Table 25	Total deaths and DALYsa from second-hand smoke, by population group and subregion, 2004.....	39
Table 26	Effects of varying assumptions on estimates of mortality caused by second-hand smoke, 2004 ^a	43
Table 27	National estimates of deaths per year attributable to SHS.....	45

FIGURES

Figure 1	Children’s SHS exposure and parental smoking, models used and survey data	8
Figure 2	Women’s SHS exposure and male active smoking in developing countries	11
Figure 3	SHS exposure and male active smoking in developed countries.....	11
Figure 4	Distribution of total deaths and DALYsa attributable to SHS, 2004.....	40

Acknowledgements

We are grateful for the thoughtful comments received from the reviewers during the preparation of this report: Stan Glantz of the University of California; Jon Samet of the John Hopkins Bloomberg School of Public Health; and Roberto Bertollini and Daniel Ferrante of the World Health Organization. We also appreciate the support received from Douglas Bettcher, Maria Neira and Luminita Sanda of the World Health Organization, and from Diana Hopkins, external consultant, who edited the report.

This report also benefited from the support of Leda Nemer and Christian Schweizer of the WHO European Centre for Environment and Health in Rome, and of the Swedish Board of Health and Welfare in the development of the method underlying this assessment, which is published separately.

This publication is the result of a joint project between the Tobacco Free Initiative (TFI) and the Public Health and the Environment Department (PHE) of the World Health Organization.

List of abbreviations

CDC	Centres for Disease Control and Prevention
CI	Confidence interval
Cal-EPA	Californian Environmental Protection Agency
DALY	Disability-adjusted life years
ECRHS	European Community Respiratory Health Survey
GYTS	Global Youth Tobacco Survey
GBD	Global burden of disease
IARC	International Agency for Research on Cancer
IHD	Ischaemic heart disease
LRI	Lower respiratory infection
OR	Odds ratio
PAF	Population attributable fraction
RR	Relative risk
SHS	Second-hand smoke
TFI	Tobacco Free Initiative
WHO	World Health Organization

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_28955

