# WHO/FAO/OIE Guidelines for the surveillance, prevention and control of taeniosis/cysticercosis

**Editor:** 

K.D. Murrell

**Associate Editors:** 

P. Dorny

A. Flisser

S. Geerts

N.C. Kyvsgaard

D. McManus

T. Nash

Z. Pawlowski

- Etiology
- Taeniosis in humans
- Cysticercosis in animals and humans
- Biology and systematics
- Epidemiology and geographical distribution
- Diagnosis and treatment in humans
- Detection in cattle and swine
- Surveillance
- Prevention
- Control
- Methods

All OIE (World Organisation for Animal Health) publications are protected by international copyright law. Extracts may be copied, reproduced, translated, adapted or published in journals, documents, books, electronic media and any other medium destined for the public, for information, educational or commercial purposes, provided prior written permission has been granted by the OIE.

The designations and denominations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the OIE concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers and boundaries.

The views expressed in signed articles are solely the responsibility of the authors. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by the OIE in preference to others of a similar nature that are not mentioned.

The designations employed and the presentation of material in this publication do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations, the World Health Organization or the World Organization for Animal Health concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The designations 'developed' and 'developing' economies are intended for statistical convenience and do not necessarily express a judgement about the stage reached by a particular country, territory or area in the development process.

The views expressed herein are those of the authors and do not necessarily represent those of the Food and Agriculture Organization of the United Nations and the World Health Organization

The views expressed in signed articles are solely the responsibility of the authors. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO or WHO in preference to others of a similar nature that are not mentioned.

#### © Copyright:

OIE (World Organisation for Animal Health), WHO (World Health Organization ) and FAO (Food and Agriculture Organization ), 2005

OIE, 12, rue de Prony, 75017 Paris, France

Tel.: 33-(0)1 44 15 18 88 Fax: 33-(0)1 42 67 09 87 http://www.oie.int

ISBN: 92-9044-656-0

#### Cover

Design: © Coup de Grace Designs, Tartu, Estonia

Conception: pbpOIE

#### **FOREWORD**

Taeniosis and cysticercosis are two parasitic diseases that in the past have not always been recognized for their importance. However, it is becoming increasingly clear that greater priority should be given to these zoonoses because of their economic impact, particularly in resource-poor countries, and their public health burden. They are also now recognized as an increasing problem in some regions, such as Africa. The now accepted linkage between epilepsy and neurocysticercosis in countries endemic for *Taenia solium* is further impetus for allocating more effort to the control of taeniosis/cysticercosis.

As is the case for all zoonoses, the control of taeniosis/cysticercosis, requires a very close collaboration between both veterinary and medical public health services at a national level. It was with the aim of assisting those responsible for taeniosis/cysticercosis control and prevention that these Guidelines were prepared and jointly published by the Food and Agriculture Organization (FAO), the World Health Organization (WHO) and the World Organisation for Animal Health (OIE). This joint publication is another example of the three organizations working together to assist their member countries in designing, implementing and standardizing control strategies against zoonoses on both national and international levels. This is the result of years of effort by FAO, WHO and OIE to educate and to organize the surveillance and control of parasitic diseases world-wide.

This FAO/WHO/OIE Guidelines for the Surveillance, Prevention and Control of Taenisosis and Cysticercosis is a compilation of the knowledge and valuable expertise of a great many internationally recognized experts on these zoonoses, accumulated over more than a hundred years of research. Without this knowledge base, effective and proven recommendations for diagnosis, treatment, prevention and control would not be possible. We would like to express our profound gratitude to these experts for their contributions and for those able to share their knowledge and advice with the authors. It is our hope that this book achieves the success it deserves.

Carlos Eddi FAO Food and Agriculture Organization

Francois-Xavier Meslin WHO World Health Organization Bernard Vallat OIE World Organisation for Animal Health

I

#### **AUTHORS**

J.C. **Allan**, Pfizer Animal Health, Licensing and Business Development, Pfizer Ltd, Sandwich, United Kingdom, E-Mail: James\_Allan@Sandwich.Pfizer.Com.

Guillermina **Avila**, Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de México, México 04510 DF, México, E-Mail: Guilleavila2000@Yahoo.Com.

J. **Brandt**, Department of Animal Health, Prince Leopold Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium, E-Mail: Jbrandt@Itg.Be.

Dolores **Correa**, Subdirección de Medicina Experimental, Instituto Nacional de Pediatría, Secretaria de Salud, México 04530 DF, México, E-Mail: mariadol@yahoo. com.

- Oscar H. **Del Brutto**, Department of Neurological Sciences, Hospital-Clínica Kennedy, Guayaquil, Ecuador.
- P. **Dorny**, Department of Animal Health, Prince Leopold Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium, E-Mail: Pdorny@Itg.Be.

Ana **Flisser**, Departamento de Microbiología y Parasitología, Facultad de Medicina, Universidad Nacional Autónoma de México, México 04510 DF, México and Dirección de Investigación. Hospital General 'Dr Manuel Gea González', Secretaria de Salud, México 14000 DF, México. E-Mail: Flisser@Servidor.Unam.Mx.

- Hector H. **Garcia**, Cysticercosis Unit, Instituto Nacional de Ciencias Neurolo´gicas, Departments of Microbiology and Pathology, Universidad Peruana, Cayetano Heredia, Peru.
- S. **Geerts**, Department of Animal Health, Prince Leopold Institute of Tropical Medicine, Nationalestraat 155, 2000 Antwerpen, Belgium, E-Mail: Sgeerts@Itg.Be.
- A. **Ito**, Department of Parasitology, Asahikawa Medical College, Asahikawa, 078-8510, Japan. E-Mail: Akira Ito [Akiraito@Asahikawa-Med.Ac.Jp].
- Niels C. **Kyvsgaard**, WHO/FAO Collaborating Center for Emerging Parasitic Zoonoses, Danish Center for Experimental Parasitology, Royal Veterinary and Agricultural University, 1870 Frederiksberg C, Denmark, E-Mail: Nck@Kvl.Dk.
- Pablo **Maravilla,** Dirección de Investigación. Hospital General 'Dr Manuel Gea González', Secretaria de Salud, México 14000 DF, México E-Mail: Maravillap@Yahoo.Com.
- D. P. **McManus**, Molecular Parasitology Laboratory, Australian Centre for International and Tropical Health and Nutrition, The Queensland Institute of Medical Research and The University of Queensland, Brisbane, Queensland 4029, Australia, E-Mail: Donm@Qimr.Edu.Au.
- H. **Meinardi**, retired Professor of epileptology Nijmegen University, The Netherlands; Past-President International Bureau for Epilepsy and International League Against Epilepsy, Netherlands, E-Mail: Meinardi@Wxs.Nl.
- K.D. **Murrell**, Danish Centre for Experimental Parasitology, WHO/FAO Collaborating Centre for Emerging Parasitic Zoonoses, Institute of Veterinary Pathobiology, Royal Veterinary and Agricultural University, 1870 Frederiksberg, Denmark, E-Mail: Kdmurrell@Comcast.Net.

Theodore E. **Nash**, Laboratory of Parasitic Diseases, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, Maryland.

Z.S. **Pawłowski**, Clinic of Tropical and Parasitic Diseases, University of Medical Sciences, Ul.Przybyszewskiego 49, 60-355 POZNAN, Poland, E-Mail: Zpawlows@Am.Poznan.Pl.

Vedantam Rajshekhar, Department of Neurological Sciences, Christian Medical College and Hospital, Vellore, India.

### **CONTENTS**

Foreword	I
AUTHORS	III
CONTENTS	V
INTRODUCTION	XI
CHAPTER 1: BIOLOGY OF TAENIA SOLIUM, TAENIA SAGINATA AND TAENIA	1
SAGINATA ASIATICA	
1. Introduction	
1.1. Morphology	
1.3. Systematics	
1.4. Host immune responses	
1. 1. 1103t inimitatic responses	
CHAPTER 2: CLINICAL CYSTICERCOSIS: DIAGNOSIS AND TREATMENT	11
2. Introduction	11
2.2. Natural history and disease	
2.2.1. Correlation of imaging and pathology	12
2.2.2. Chronic (inactive) cysticercosis associated with brain calcifications	13
2.2.3. Patterns of presentation and type of involvement	
2.2.3.1. Solitary cysticercus granuloma	
2.2.3.2. Extraparenchymal neurocysticercosis	14
2.2.3.3. Massive infection	
2.2.3.4. Involvement of other organs	
2.3.1. Overall assessment and available diagnostic tools	
2.3.2. Diagnostic criteria for neurocysticercosis	
2.3.2.1. Introduction	
2.3.2.2. Definition of diagnostic criteria	
2.3.2.3. Degrees of diagnostic certainty	
2.4. Treatment of neurocysticercosis	20
2.4.1. Introduction	20
2.4.2. Symptomatic medication.	21
2.4.3. Anthelminthic medications-indications and use	22
2.4.4. Management in field conditions	24
2.4.5. Screening and treatment of tapeworms	24
CHAPTER 3: EPIDEMIOLOGY OF TAENIOSIS AND CYSTICERCOSIS	27
3. Introduction	27
3.1. Global distribution of <i>Taenia solium</i> taeniosis/cysticercosis	27
3.1.1. Africa	27
3.1.2. Latin America	29
3.1.3. Europe	30
3.1.4. Asia	
3.1.5. United States of America: The impact of immigration and travel	32
3.2. Global distribution and occurrence of taeniosis/cysticercosis due to Taenia saginata	33

3.2.1. Taeniosis	33
3.2.2. Bovine cysticercosis	33
3.3. Global distribution and occurrence of Taenia saginata asiatica (Asian Taenia)	
3.4. Factors affecting transmission of taeniid cestodes	33
3.4.1. Taenia solium	34
3.4.1.1. Transmission from pigs to humans	34
3.4.1.2. Transmission from humans to pigs	34
3.4.1.3. Person to person transmission.	35
3.4.2. Taenia saginata	
3.4.2.1. Transmission from cattle to humans	
3.4.2.2. Transmission from humans to cattle	
3.4.3. Extrinsic factors affecting taeniid egg survival and dispersal	
3.4.3.1. Egg output	
3.4.3.2. Egg dispersal agents	
3.4.3.3. Factors affecting egg viability and survival	
3.4.4. Intrinsic factors affecting transmission.	
3.4.4.1. Immunological mechanisms	
3.4.5. The role of livestock management in transmission	
3.5. Summary of risk factors for taeniosis and cysticercosis	
3.5.1. Taenia solium	
3.5.2. Taenia saginata	43
3.5.3. Taenia saginata asiatica	43
CHAPTER 4: DETECTION AND DIAGNOSIS	45
4. Introduction	45
4.1. Diagnosis of taeniosis	45
4.1.1. Diagnosis based on morphological criteria	
4.1.2. Differentiation based on enzyme electrophoresis	
4.1.3. Differentiation based on molecular techniques	
4.1.4. Questioning of tapeworm carriers	
4.1.5. Coprological examinations	
4.1.5.1. Conventional fecal examinations	
4.1.5.2. Peri-anal swabs	
4.1.5.3. Coproantigen detection	47
4.1.5.4. Copro-PCR	47
4.1.6. Serological tests	47

## 预览已结束,完整报告链接和二维码如下:

https://www.yunbaogao.cn/report/index/report?reportId=5\_29879



