

BURULI ULCER

Mycobacterium ulcerans infection



Edited by

Dr Kingsley Asiedu
Dr Robert Scherpbier
Dr Mario Raviglione





Buruli ulcer commonly affects poor people who live near rivers or wetlands.
(Photo: Augustin Guedenon)

**This document was published with contributions from
the Association Française Raoul Follereau, Paris, France
and The Nippon Foundation, Tokyo, Japan.**

WHO/CDS/CPE/GBUI/2000.1
Distr.: General
Original : English

Buruli ulcer

Mycobacterium ulcerans infection

Editors :

Dr Kingsley Asiedu

Dr Robert Scherpbier

Dr Mario Raviglione



World Health Organization

Global Buruli Ulcer Initiative

Copyright © World Health Organization (2000)

This document is not a formal publication of the World Health Organization (WHO) and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, reproduced, or translated in part, but not for sale or for use in conjunction with commercial purposes.

The views expressed herein by named authors are solely the responsibility of those authors.

List of contributors

Dr Pius Agbenorku, Department of Plastic Surgery, School of Medical Sciences, Kumasi, **Ghana**. Dr George Amofah, Public Health Division, Ministry of Health, Accra, **Ghana**. Dr Kingsley Asiedu, Communicable Diseases, World Health Organization, Geneva, **Switzerland**. Mr Karim Chemlal, Department of Microbiology, Institute of Tropical Medicine, Antwerp, **Belgium**. Dr Karen Dobos, Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, Georgia, **USA**. Dr Samuel Etuaful, St Martin's Catholic Hospital, Agroyesum, **Ghana**. Dr Mark Evans, Division of Infectious Diseases, St George's Hospital Medical School, London, **United Kingdom**. Dr Kathleen George, Cell Biology, Rocky Mountain Laboratories, Hamilton, Montana, **USA**. Professor Jacques Grosset, Bactériologie et Hygiène, Faculté de Médecine, Pitié-Salpêtrière, Paris, **France**. Dr Augustin Guedenon, Dermato-Vénérologue, Cotonou, **Benin**. Dr John Hayman, Department of Pathology, Box Hill Hospital, Victoria, **Australia**. Dr Paul Johnson, Department of Infectious Diseases, Monash Medical Center, Melbourne, **Australia**. Professor Jean-Marie Kanga, Dermatology Centre, Treichville University Hospital, Abidjan, **Côte d'Ivoire**. Dr C. Harold King, Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, Georgia, **USA**. Dr Wayne Meyers, Armed Forces Institute of Pathology, Washington, DC, **USA**. Dr Eugene Nyarko, WHO Regional Office for Africa, Harare, **Zimbabwe**. Professor Françoise Portaels, Department of Microbiology, Institute of Tropical Medicine, Antwerp, **Belgium**. Dr Mario Raviglione, Communicable Diseases, World Health Organization, Geneva, **Switzerland**. Dr Robert Scherpbier, Communicable Diseases, World Health Organization, Geneva, **Switzerland**. Dr Pam Small, Cell Biology, Rocky Mountain Laboratories, Hamilton, Montana, **USA**. Dr Tim Stinear, Department of Microbiology, Monash University, Melbourne, **Australia**. Dr Jordan Tappero, Centers for Disease Control and Prevention, Atlanta, Georgia, **USA**. Dr Napo Tignokpa, Dermatology Centre, Leprosy Programme, Lomé, **Togo**. Dr Mark Wansbrough-Jones, Division of Infectious Diseases, St George's Hospital Medical School, London, **United Kingdom**.

Acknowledgement

The Global Buruli Ulcer Initiative receives funding from The Nippon Foundation, Tokyo, Japan. This project could not have succeeded without the direct support to the endemic countries by nongovernmental organizations, the political commitments from leaders of the affected countries and the cooperation of their ministries of health. The comments of Frank Theakston, WHO European Office, Copenhagen, Denmark are greatly appreciated. Many thanks to those who contributed to this monograph and those who provided photographs. The secretarial support of Stéphanie Jourdan, Sylvie Lamy Quique and Jeanette Dadzie is recognized.

Contents

Chapters

Executive Summary	1
Chapter 1. Introduction	5
Chapter 2. Epidemiology	8
Chapter 3. Microbiology	13
Chapter 4. Immunology	17
Chapter 5. Modern Diagnostic Techniques	21
Chapter 6. Toxin	29
Chapter 7. Pathology	33
Chapter 8. Clinical Features and Treatment	35
Chapter 9. Prevention, Surveillance and Control	47
Chapter 10. Economic and Social Impact	55
Chapter 11. Research Coordination	59

Tables

Table 1.	Regions and countries with reported cases of Buruli ulcer worldwide (unpublished)
Table 2.	Fingerprinting techniques used for <i>M. ulcerans</i> (unpublished, Françoise Portaels)
Table 3.	Organization of laboratory services (Yamoussoukro Report)
Table 4.	Proposed clinical classification (CC) for the management of Buruli ulcer
Table 5.	Guidelines for the clinical management of Buruli ulcer (Yamoussoukro Report)
Table 6.	Development of national control programmes (unpublished)

Figures

- Fig. 1. Map of Africa showing the geographical transition of Buruli ulcer before and after 1980 (Françoise Portael)
- Fig. 2. Geographical distribution of Buruli ulcer cases worldwide
- Fig. 3. Serological diagnosis of Buruli ulcer: preliminary results
- Fig. 4. Structure of *M. ulcerans* toxin (published in Science, 1999)
- Fig. 5. Distribution of treatment costs (published in American Journal of Tropical Medicine and Hygiene, 1998)
- Fig. A5.1a. Trends of Buruli ulcer cases in Côte d'Ivoire, 1978–1997
- Fig. A5.1b,c. Geographical distribution of Buruli ulcer cases in Côte d'Ivoire, 1978 and 1997
- Fig. A5.2a. Trends of Buruli ulcer cases in Benin, 1989–1998
- Fig. A5.2b,c. Geographical distribution of Buruli ulcer cases in Benin 1989 and 1997

Plates

- Plate 1. Ziehl-Neelsen stained smear from a Buruli ulcer showing extracellular acid-fast bacilli (Photo: Wayne Meyers)
- Plate 2. Ghost cells (Photo: Wayne Meyers)
- Plate 3. Ghost cells and vasculitis (Photo: Wayne Meyers)
- Plate 4. Biopsy specimen from the edge of an ulcer showing undermining of the dermis and massive necrosis of the skin, dermis, subcutis and the fascia (Photo: Wayne Meyers)
- Plate 5. Undermined edges of Buruli ulcer. Note the characteristic yellowish-white necrotic base (cotton wool-like appearance) (Photo: John Hayman)
- Plate 6. Lymphadenopathy in Buruli ulcer. The parenchyma of the node is necrotic and the capsule is heavily infiltrated by acid-fast bacilli. (Photo: Wayne Meyers)
- Plate 7. X-ray of the foot showing destruction of the bone. Note the patient had Buruli ulcer on the dorsum of the foot (Photo: Wayne Meyers & Battista Priuli)
- Plate 8. A papule (Photo: John Hayman)

- Plate 9. A nodule (Photo: Mark Evans)
- Plate 10. Plaque (Photo: Mark Evans)
- Plate 11. Plaque (Photo: Kingsley Asiedu)
- Plate 12. Nonulcerative oedema (Photo: Samuel Etuaful)
- Plate 13. Ulcerative oedema (Photo: Kingsley Asiedu)
- Plate 14. Ulcerative lesion (Photo: John Hayman)
- Plate 15. Ulcerative lesion (Photo: Roger Pradinaud)
- Plate 16. Ulcerative lesion (Photo: Augustin Guedenon)
- Plate 17. Ulcerative lesion (Photo: Roger Pradinaud)
- Plate 18. Ulcerative lesion (Photo: Mark Evans)
- Plate 19. Ulcerative lesion (Photo: Mark Evans)
- Plate 20. Ulcerative lesion (Photo: John Hayman)
- Plate 21. Ulcerative lesion (Photo: Kingsley Asiedu)
- Plate 22. Ulcerative lesion (Photo: Marco Pirovano)
- Plate 23. Ulcerative lesion (Photo: Marco Pirovano)
- Plate 24. Ulcerative (Photo: John Hayman)
- Plate 25. Ulcerative lesion (Photo: National Buruli Ulcer Control Programme, Côte d'Ivoire)
- Plate 26. Postulcerative sequelae (Photo: Augustin Guedenon)

预览已结束，完整报告链接和二维码如下：

https://www.yunbaogao.cn/report/index/report?reportId=5_30499

