

# ***Indicators for assessing Vitamin A Deficiency and their application in monitoring and evaluating intervention programmes***



World Health  
Organization

**INDICATORS FOR ASSESSING  
VITAMIN A DEFICIENCY  
AND THEIR APPLICATION IN  
MONITORING AND EVALUATING  
INTERVENTION PROGRAMMES**



World Health Organization

---

Reprinted in 1998 with the following adjustments:

Paragraphs renumbered from paragraph 29 onwards

Table 7: minimum sample size for 0.01 prevalence, from 153 650 to 153 640

Table 8: minimum sample size for  $\geq 1\%$  -  $<5\%$ , from 4706 and 753 to 9508 and 1522

Annex 1: updated

---

© World Health Organization 1996

This document is not a formal publication of the World Health Organization (WHO), and all rights are reserved by the Organization. The document may, however, be freely reviewed, abstracted, quoted, reproduced or translated, in part or in whole, but not for sale or for use in conjunction with commercial purposes.

The views expressed in documents by named authors are solely the responsibility of those authors.

## ACKNOWLEDGMENTS

WHO wishes to thank the Task Force "Sight and Life" for the financial support that made the production of this document possible; UNICEF for its financial contribution towards convening a technical consultation on the subject in 1992; and the many who took time to read and comment on a review version of this document whose suggestions are reflected herein. Grateful acknowledgment is also due to Eileen Brown for her tireless efforts in revising the manuscript, and to James Akre for his editorial assistance; both are from WHO's Programme of Nutrition, Geneva.



# TABLE OF CONTENTS

Acknowledgments .....	iii
List of tables .....	vii
List of annexes .....	viii
Abbreviations .....	viii
INTRODUCTION .....	i
Background .....	i
PART I. SUMMARY OF INDICATORS FOR VAD SURVEILLANCE .....	5
1.1 Assessing VAD prevalence .....	5
1.2 Identifying high-risk areas/populations for interventions .....	8
1.3 Measuring progress towards long-term micronutrient goals .....	12
1.4 Monitoring and evaluating VAD control programmes .....	14
PART II. USING INDICATORS FOR VAD SURVEILLANCE .....	16
2.1 Purposes of VAD surveillance .....	16
2.2 Factors in selecting groups to measure .....	16
2.3 Appropriate groups to measure .....	17
2.4 Selecting indicators .....	19
2.5 Availability and interpretation of reference data .....	20
2.6 Presentation and interpretation of survey results .....	20
2.7 Definition of a public health problem .....	21
PART III. BIOLOGICAL INDICATORS OF VITAMIN A STATUS .....	22
3.1 Clinical indicators: Xerophthalmia .....	22
3.2 Subclinical indicators .....	24
3.2.1 Functional indicator: Night blindness .....	24
3.2.2 Biochemical indicators .....	26
Serum retinol .....	26
Breast milk vitamin A concentration .....	30
Relative dose response tests (RDR and MRDR) .....	34
Serum 30-day dose response test (+S30DR) .....	38
3.2.3 Histological indicator: Impression cytology .....	39
PART IV. ECOLOGICAL AND RELATED INDICATORS ASSOCIATED WITH RISK OF VAD .....	45
4.1 Purpose and use of indirect indicators associated with VAD .....	45
4.2 Nutritional status and diet-related indicators .....	46

4.2.1	Breast-feeding patterns	46
4.2.2	Anthropometric indicators of PEM	46
4.2.3	Prevalence of low birth weight (LBW)	47
4.2.4	Market and household food availability	47
4.2.5	Dietary patterns of vulnerable groups	48
4.2.6	Semi-quantitative and qualitative measures of food consumption	49
4.2.7	Beliefs and attitudes concerning food	50
4.3	Illness-related indicators	50
4.3.1	Immunization coverage rates	51
4.3.2	Measles case fatality rate	51
4.3.3	Disease prevalence rates	51
4.4	Socioeconomic indicators	53
4.4.1	Maternal education and literacy	53
4.4.2	Income/employment	53
4.4.3	Water supply and sanitation	54
4.4.4	Access to health and social services	54
4.4.5	Access to land	55
4.4.6	Access to agricultural services and inputs	55
PART V.	INTEGRATED APPROACHES TO CONTROLLING VAD	58
5.1	Improving dietary quality and quantity through dietary diversification	58
5.2	Improving dietary quality through food fortification	59
5.3	Supplementation	59
5.4	Public health measures	60

## List of Tables

1. Indicators of clinical vitamin A deficiency - xerophthalmia - in children 6-71 months of age .....	6
2. Biological indicators of subclinical vitamin A deficiency in children 6-71 months of age .....	7
3. Types of indicators useful for achieving broad surveillance objectives .....	9
4. A. Relative ranking on a population base of some biologic surveillance purposes .....	
B. Ecologic indicators of areas/populations at risk of VAD .....	
5. Core indicators for assessing progress towards the goal of VAD by the year 2000 .....	
6. Examples of process indicators for monitoring and evaluation programmes .....	
7. Classification of xerophthalmia, minimum prevalence sizes for defining clinical VAD of public health importance children under 6 years of age .....	
8. Prevalence of night blindness in children 24-71 month ages for identifying a VAD public health problem .....	
9. Prevalence of serum values of vitamin A $\leq 0.70 \mu\text{mol/l}$ minimum sample sizes for identifying a VAD public health problem .....	
10. Prevalence of breast-milk values $\leq 1.05 \mu\text{mol/l}$ ( $\leq 8 \mu\text{g}$ ) of lactating women and minimum sample sizes for identifying a VAD public health problem .....	
11. Prevalence of a positive RDR or MRDR and minimum sample sizes for identifying a VAD public health problem .....	
12. Prevalence of positive +S30DR value in children 1-6 years sample sizes for identifying a VAD public health problem .....	
13. Comparison of the sensitivity and specificity of impression cytology indicators of VAD .....	
14. Prevalence of abnormal impression cytology in children minimum sample size for identifying a VAD public health problem .....	
15. Ecological indicators and their potential uses in VAD surveillance .....	



预览已结束，完整报告链接和二维码如下：

[https://www.yunbaogao.cn/report/index/report?reportId=5\\_30654](https://www.yunbaogao.cn/report/index/report?reportId=5_30654)